

CRPL-F118

Library, N. W. Bldg.

FOR OFFICIAL USE

Reference book not to be
taken from the Library.

IONOSPHERIC DATA

ISSUED
JUNE 1954

U. S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
CENTRAL RADIO PROPAGATION LABORATORY
BOULDER, COLORADO

IONOSPHERIC DATA

CONTENTS

	<u>Page</u>
Symbols, Terminology, Conventions	2
World-Wide Sources of Ionospheric Data	5
Hourly Ionospheric Data at Washington, D. C..	7, 12, 24, 49
Ionospheric Storminess at Washington, D. C. .	7, 33
Radio Propagation Quality Figures	8, 37
Observations of the Solar Corona	9, 40
Relative Sunspot Numbers	10, 44
Observations of Solar Flares	10, 46
Indices of Geomagnetic Activity	11, 47
Sudden Ionosphere Disturbances	11, 48
Erratum	11
Tables of Ionospheric Data	12
Graphs of Ionospheric Data	49
Index of Tables and Graphs of Ionospheric Data in CRPL-F118	85

SYMBOLS, TERMINOLOGY, CONVENTIONS

Beginning with data reported for January 1952, the symbols, terminology, and conventions for the determination of median values used in this report (CRPL-F series) conform as far as practicable to those adopted at the Sixth Meeting of the International Radio Consultative Committee (C.C.I.R.) in Geneva, 1951. Excerpts concerning symbols and terminology from Document No. 626-E of this Meeting are given on pages 2-7 of the report CRPL-F89, "Ionospheric Data," issued January 1952. Reprints of these pages are available upon request.

Beginning with data for January 1945, median values are published wherever possible. Where averages are reported, they are, at any hour, the average for all the days during the month for which numerical data exist.

The following conventions are used in determining the medians for hours when no measured values are given because of equipment limitations and ionospheric irregularities. Symbols used are those given in Document No. 626-E referred to above.

a. For all ionospheric characteristics:

Values missing because of A, C, F, L, M, N, Q, S, or T are omitted from the median count.

b. For critical frequencies and virtual heights:

Values of f_oF_2 (and f_oE near sunrise and sunset) missing because of E are counted as equal to or less than the lower limit of the recorder. Values of $h'F_2$ (and $h'E$ near sunrise and sunset) missing for this reason are counted usually as equal to or greater than the median. Other characteristics missing because of E are omitted from the median count.

Values missing because of D are counted as equal to or greater than the upper limit of the recorder.

Values missing because of G are counted:

1. For f_oF_2 , as equal to or less than f_oF_1 .
2. For $h'F_2$, as equal to or greater than the median.

The symbol W is included in the median count only when it replaces a height characteristic. This practice represents a change from that listed in issues previous to CRPL-F78.

Values missing for any other reason are omitted from the median count.

c. For MUF factor (M-factors):

Values missing because of G or W are counted as equal to or less than the median.

Values missing for any other reason are omitted from the median count.

d. For sporadic E (Es):

Values of fEs missing because of E or G (and B when applied to the daytime E region only) are counted as equal to or less than the median foE, or equal to or less than the lower frequency limit of the recorder.

Values of fEs missing for any other reason, and values of h'Es missing for any reason at all are omitted from the median count.

Beginning with data for November 1945, doubtful monthly median values for ionospheric observations at Washington, D. C., are indicated by parentheses, in accordance with the practice already in use for doubtful hourly values. The following are the conventions used to determine whether or not a median value is doubtful:

1. If only four values or less are available, the data are considered insufficient and no median value is computed.

2. For the F2 layer, if only five to nine values are available, the median is considered doubtful. The E and F1 layers are so regular in their characteristics that, as long as there are at least five values, the median is not considered doubtful.

3. For all layers, if more than half of the values used to compute the median are doubtful (either doubtful or interpolated), the median is considered doubtful.

The same conventions are used by the CRPL in computing the medians from tabulations of daily and hourly data for stations other than Washington, beginning with the tables in IRPL-F18.

The tables and graphs of ionospheric data are correct for the values reported to the CRPL, but, because of variations in practice in the interpretation of records and scaling and manner of reporting of values, may at times give an erroneous conception of typical ionospheric characteristics at the station. Some of the errors are due to:

- a. Differences in scaling records when spread echoes are present.
- b. Omission of values when f_oF_2 is less than or equal to f_oF_1 , leading to erroneously high values of monthly averages or median values.
- c. Omission of values when critical frequencies are less than the lower frequency limit of the recorder, also leading to erroneously high values of monthly average or median values.

These effects were discussed on pages 6 and 7 of the previous F-series report IRPL-F5.

Ordinarily, a blank space in the fEs column of a table is the result of the fact that a majority of the readings for the month are below the lower limit of the recorder or less than the corresponding values of f_oE . Blank spaces at the beginning and end of columns of $h'F_1$, f_oF_1 , $h'E$, and f_oE are usually the result of diurnal variation in these characteristics. Complete absence of medians of $h'F_1$ and f_oF_1 is usually the result of seasonal effects.

The dashed-line prediction curves of the graphs of ionospheric data are obtained from the predicted zero-muf contour charts of the CRPL-D series publications. The following points are worthy of note:

- a. Predictions for individual stations used to construct the charts may be more accurate than the values read from the charts since some smoothing of the contours is necessary to allow for the longitude effect within a zone. Thus, inasmuch as the predicted contours are for the center of each zone, part of the discrepancy between the predicted and observed values as given in the F series may be caused by the fact that the station is not centrally located within the zone.
- b. The final presentation of the predictions is dependent upon the latest available ionospheric and radio propagation data, as well as upon predicted sunspot number.

- c. There is no indication on the graphs of the relative reliability of the data; it is necessary to consult the tables for such information.

The following predicted smoothed 12-month running-average Zürich sunspot numbers were used in constructing the contour charts:

Month	Predicted Sunspot Number									
	1954	1953	1952	1951	1950	1949	1948	1947	1946	1945
December		15	33	53	86	108	114	126	85	38
November		16	38	52	87	112	115	124	83	36
October		17	43	52	90	114	116	119	81	23
September		18	46	54	91	115	117	121	79	22
August		18	49	57	96	111	123	122	77	20
July		20	51	60	101	108	125	116	73	
June		21	52	63	103	108	129	112	67	
May	10	22	52	68	102	108	130	109	67	
April	10	24	52	74	101	109	133	107	62	
March	11	27	52	78	103	111	133	105	51	
February	12	29	51	82	103	113	133	90	46	
January	14	30	53	85	105	112	130	88	42	

WORLD - WIDE SOURCES OF IONOSPHERIC DATA

The ionospheric data given here in tables 1 to 72 and figures 1 to 144 were assembled by the Central Radio Propagation Laboratory for analysis and correlation, incidental to CRPL prediction of radio propagation conditions. The data are median values unless otherwise indicated. The following are the sources of the data in this issue:

University of Graz:
Graz, Austria

Meteorological Service of the Belgian Congo and Ruanda-Urundi:
Leopoldville, Belgian Congo

University of Sao Paulo:
Sao Paulo, Brazil

British Department of Scientific and Industrial Research, Radio
Research Board:
Falkland Is.
Inverness, Scotland
Khartoum, Sudan (University College of Khartoum)
Port Lockroy
Singapore, British Malaya
Slough, England

Defence Research Board, Canada:

Baker Lake, Canada
 Churchill, Canada
 Fort Chimo, Canada
 Ottawa, Canada
 Prince Rupert, Canada
 Resolute Bay, Canada
 Winnipeg, Canada

Radio Wave Research Laboratories, National Taiwan University, Taipei,
 Formosa, China:

Formosa, China

Icelandic Post and Telegraph Administration:

Reykjavik, Iceland

All India Radio (Government of India), New Delhi, India:

Bombay, India
 Delhi, India
 Madras, India
 Tiruchy (Tiruchirapalli), India

Indian Council of Scientific and Industrial Research, Radio Research
 Committee:

Calcutta, India

Ministry of Postal Services, Radio Research Laboratories, Tokyo, Japan:

Akita, Japan
 Tokyo (Kokubunji), Japan
 Wakkanai, Japan
 Yamagawa, Japan

Christchurch Geophysical Observatory, New Zealand Department of Scientific
 and Industrial Research:

Rarotonga, Cook Is.

South African Council for Scientific and Industrial Research:

Capetown, Union of South Africa
 Johannesburg, Union of South Africa
 Nairobi, Kenya (East African Meteorological Department)

Research Laboratory of Electronics, Chalmers University of Technology,
 Gothenburg, Sweden:

Kiruna, Sweden

United States Army Signal Corps:

Adak, Alaska
 White Sands, New Mexico

National Bureau of Standards (Central Radio Propagation Laboratory):
Anchorage, Alaska
Fairbanks, Alaska (Geophysical Institute of the University
of Alaska)
Guam I.
Maui, Hawaii
Puerto Rico, W. I.
San Francisco, California (Stanford University)
Washington, D. C.

HOURLY IONOSPHERIC DATA AT WASHINGTON, D. C.

The data given in tables 73 through 84 follow the scaling practices given in the report IRPL-C61, "Report of International Radio Propagation Conference," pages 36 to 39, and the median values are determined by the conventions given above under "Symbols, Terminology, Conventions." Beginning with September 1949, the data are taken at Ft. Belvoir, Virginia.

IONOSPHERIC STORMINESS AT WASHINGTON, D.C.

Table 85 presents ionosphere character figures for Washington, D. C., during May 1954, as determined by the criteria given in the report IRPL-R5, "Criteria for Ionospheric Storminess," together with Cheltenham, Maryland, geomagnetic K-figures, which are usually covariant with them.

See the last column of table 87a in this issue for the April 1954 character figures for Washington, D. C., which were not published in last month's issue.

RADIO PROPAGATION QUALITY FIGURES

Tables 87a and 87b give for April 1954 the radio propagation quality figures for the North Atlantic area, the relevant CRPL advance and short-term forecasts, a summary geomagnetic activity index and sundry comparisons, specifically as follows:

- (a) radio propagation quality figures, Q_a , separately for each 6-hour interval of the Greenwich day, viz., 00-06, 06-12, 12-18, 18-24 hours UT (Universal Time or GCT).
- (b) whole-day radio quality indices (beginning October 1952). Each index is a weighted average of the four quarter-day Q_a -figures, before rounding off, with half weight given to quality grades 5 and 6. This procedure tends to give whole-day indices suitable for comparison with whole-day advance forecasts which designate whenever possible the days when significant disturbance or unusually quiet conditions will occur.
- (c) short-term forecasts, issued by CRPL every six hours (nominally one hour before 00^h, 06^h, 12^h, 18^h UT) and applicable to the period 1 to 13 (especially 1 to 7) hours ahead. Note that new scoring rules have been adopted beginning with October 1952 data.
- (d) advance forecasts, issued semiweekly (CRPL-J reports) and applicable 1 to 3 or 4 days ahead, 4 or 5 to 7 days ahead, and 8 to 25 days ahead. These forecasts are scored against the whole-day quality indices.
- (e) half-day averages of the geomagnetic K indices measured by the Cheltenham Magnetic Observatory of the U. S. Coast and Geodetic Survey.
- (f) illustration of the comparison of short-term forecasts with Q_a -figures and also with estimates of radio quality based on CRPL observations only.
- (g) illustration of the outcome of advance forecasts (1 to 3 or 4 days ahead) and, for comparison, the outcome of a type of "blind" forecast. For the latter the frequency for each quality grade, as determined from the distribution of quality grades in the four most recent months of the current season, is partitioned among the grades observed in the current month in proportion to the frequencies observed in the current month.

These radio propagation quality figures, Q_a , are prepared from radio traffic data reported to CRPL by American Telephone and Telegraph Company, Mackay Radio and Telegraph Company, RCA Communications, Inc., Marconi Company, British Admiralty Signal and Radar Establishment, and the following agencies of the U. S. Government:--Coast Guard, Navy, Army Signal Corps, and U. S. Information Agency. The method of calculation, summarized below, is similar to that described in a 1946 report, IRPL-R31, now out of print. Only reports of radio transmission on North Atlantic paths closely approximating New York-London are included in the estimation of quality.

The original reports are submitted on various scales and for various time intervals. The observations for each 6-hour interval are averaged on the quality scale of the original reports. These 6-hour indices are then adjusted to the 1 to 9 quality-figure scale by a conversion table prepared by comparing the distribution of these indices for at least four months, usually a year, with a master distribution determined from analysis of the reports originally made on the 1 to 9 quality-figure scale. A report whose distribution is the same as the master is thereby converted linearly to the Q -figure scale. The 6-hourly quality figures are (subjectively) weighted means of the reports received for that period. These 6-hourly quality figures replace, beginning January 1953, the half-daily quality figures which formerly appeared in this table. (These forecasts and quality indices are prepared by the North Atlantic Radio Warning Service, the CRPL forecasting center at Ft. Belvoir, Virginia.)

Table 86 gives for April 1954, the radio propagation quality figures for the North Pacific area, the relevant CRPL advance and short-term forecasts, and sundry comparisons, specifically as follows:

- (a) radio propagation quality figures, Q_p , separately for each of three 9-hour intervals of the Greenwich day, viz., 03-12, 09-18 and 18-03 UT (Universal Time or GCT).
- (b) whole-day radio quality indices for each Greenwich day. These are derived from the same basic data as the 9-hour indices, separately reduced.
- (c) short-term forecasts, issued daily at 02, 09 and 18 hours UT.
- (d) advance forecasts, issued semiweekly (CRPL-Jp reports) and applicable 1 to 3 or 4 days ahead, 4 or 5 to 7 days ahead, and 8 to 25 days ahead. These forecasts are scored against the whole day quality indices.

These radio quality indices, Q_p , refer to radio propagation on optimum frequencies over moderately long transmission paths in the North Pacific area. Typical paths are Anchorage (Alaska) to Seattle, or Anchorage to Tokyo. The indices are derived from reports submitted regularly by communications agencies of the U. S. Army and Air Force, and by Aeronautical Radio, Inc. The method of derivation of Q_p differs from that of Q_a . For Q_p , each reported index is converted into a deviation (usually) from the 3-monthly mean for that index, in units of the standard deviation. These deviations are averaged for all reports for a given 9-hour period. The average is then put on the 1 to 9 Q-scale with an assumed standard deviation of 1.25 and assumed means of 5.33, 5.33, and 6.00, respectively, for the 03-12, 09-18 and 18-03 periods, and 5.67 for the whole day period. (These forecasts and quality indices are prepared by the North Pacific Radio Warning Service, the CRPL forecasting center at Anchorage, Alaska.)

These quality figures are, in effect, a consensus of reported radio propagation conditions. The reasons for low quality are not necessarily known and may not be limited to ionospheric storminess. For instance, low quality may result from improper frequency usage for the path and time of day. Although, wherever it is reported, frequency usage is included in the rating of reports, it must often be an assumption that the reports refer to optimum working frequencies. It is more difficult to eliminate from the indices conditions of low quality because of multipath, interference, etc. These considerations should be taken into account in interpreting research correlations between the Q-figures and solar, auroral, geomagnetic or similar indices.

OBSERVATIONS OF THE SOLAR CORONA

Tables 88 through 90 give the observations of the solar corona during May 1954, obtained at Climax, Colorado, by the High Altitude Observatory of Harvard University and the University of Colorado. Tables 91 through 93 list the coronal observations obtained at Sacramento Peak, New Mexico, during May 1954, derived by Harvard College Observatory as a part of its performance of a research contract with the Upper Air Research Observatory, Geophysical Research Directorate, Air Force Cambridge Research Center. The data are listed separately for east and west limbs at 5-degree intervals of position angle north and south of the Solar Equator at the limb. The time of observation is given to the nearest tenth of a day, GCT.

Table 88 gives the intensities of the green (5303A) line of the emission spectrum of the solar corona; table 89 gives similarly the intensities of the first red (6374A) coronal line; and table 90, the intensities of the second red (6702A) coronal line; all observed at Climax in May 1954.

Table 91 gives the intensities of the green (5303A) coronal line; table 92, the intensities of the first red (6374A) coronal line; and table 93, the intensities of the second red (6702A) coronal line; all observed at Sacramento Peak in May 1954.

The following symbols are used in tables 88 through 93: a, observation of low weight; -, corona not visible; and X, position angle not included in plate estimates.

RELATIVE SUNSPOT NUMBERS

Table 94 lists the daily provisional Zurich relative sunspot number, R_z , for May 1954, as communicated by the Swiss Federal Observatory. Table 95 contains the daily American relative sunspot number, R_A' , for April 1954, as compiled by the Solar Division, American Association of Variable Star Observers.

OBSERVATIONS OF SOLAR FLARES

Table 96 gives the preliminary record of solar flares reported to the CRPL. These reports are communicated on a rapid schedule at the sacrifice of detailed accuracy. Definitive and complete records are published later in the Quarterly Bulletin of Solar Activity, I.A.U., in various observatory publications, and elsewhere. The present listing serves to identify and roughly describe the phenomena observed. Details should be sought from the reporting observatory.

Reporting directly to the CRPL are the following observatories: Mt. Wilson, McMath-Hulbert, U. S. Naval, Wendelstein, Kanzel and High Altitude at Sacramento Peak, New Mexico. The remainder report to Meudon (Paris) and the data are taken from the Paris-URSIGRAM broadcast, monitored fairly regularly by the CRPL. The data on solar flares reported from Sacramento Peak, New Mexico, communicated by the High Altitude Observatory at Boulder, Colorado, are provided by Harvard University as the result of work undertaken on an Air Materiel Command Research and Development Contract administered by the Air Force Cambridge Research Laboratories.

The table lists for each flare the reporting observatory, date, times of beginning and ending of observation, duration (when known), total area (corrected for foreshortening), and heliographic coordinates. For the maximum phase of the flare is given the time, intensity, area relative to the total area, and the importance. The column "SID observed" is to indicate when a sudden ionosphere disturbance, noted elsewhere in these reports, occurred at the time of a flare. Times are in Universal Time (GCT).

INDICES OF GEOMAGNETIC ACTIVITY

Table 97 lists various indices of geomagnetic activity based on data from magnetic observatories widely distributed throughout the world. The indices are: (1) preliminary international character-figures, C; (2) geomagnetic planetary three-hour-range indices, Kp; (3) magnetically selected quiet and disturbed days.

The C-figure is the arithmetic mean of the subjective classification by all observatories of each day's magnetic activity on a scale of 0 (quiet) to 2 (storm). The magnetically quiet and disturbed days are selected by the international scheme outlined on pages 219-227 in the December 1943 issue of Terrestrial Magnetism and Atmospheric Electricity. The details of the currently used method follow. For each day of a month, its geomagnetic activity is assigned by weighting equally the following three criteria: (1) the sum of the eight Kp's; (2) the greatest Kp; and (3) the sum of the squares of the eight Kp's.

Kp is the mean standardized K-index from 11 observatories between geomagnetic latitudes 47 and 63 degrees. The scale is 0 (very quiet) to 9 (extremely disturbed), expressed in thirds of a unit, e.g., 5- is $4 \frac{2}{3}$, 5o is $5 \frac{0}{3}$, and 5+ is $5 \frac{1}{3}$. This planetary index is designed to measure solar particle-radiation by its magnetic effects, specifically to meet the needs of research workers in the ionospheric field. A complete description of Kp has appeared in Bulletin 12b, "Geomagnetic Indices C and K, 1948," published in Washington, D. C., 1949, by the Association of Terrestrial Magnetism and Electricity, International Union of Geodesy and Geophysics. Kp is available from 1937 to date as noted in F108.

The Committee on Characterization of Magnetic Disturbance, ATME, IUGG, has kindly supplied this table. The Meteorological Office, De Bilt, Holland, collects the data and compiles C and selected days. The Chairman of the Committee computes the planetary index. Current tables are also published quarterly in the Journal of Geophysical Research along with data on sudden commencements (sc) and solar flare effects (sfe).

SUDDEN IONOSPHERE DISTURBANCES

Table 98 shows that no sudden ionosphere disturbances were observed at Ft. Belvoir, Virginia, during the months of April and May 1954.

ERRATUM

CRPL-F114, p. 12, table 1: At 02 in foF2 column, the value should be 2.6;
 p. 21, table 50: In 02 column, the median value should be 2.6;
 p. 44, fig. 1: At 02 in foF2 curve, reading should be 2.6.

TABLES OF IONOSPHERIC DATA

Table 1

Washington, D. C. (38.7°N, 77.1°W)

May 1954

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	280	2.9					2.5	3.1
01	290	2.5					3.1	
02	290	2.3					2.9	3.1
03	290	2.2					2.9	3.2
04	(280)	2.0					3.2	3.1
05	250	2.7					3.0	3.3
06	310	3.6	220	3.2	120	1.8	3.1	3.25
07	330	4.1	210	3.5	110	2.3	3.8	3.25
08	350	4.4	210	3.8	100	2.6	4.4	3.1
09	340	4.8	200	4.0	100	2.8	4.4	3.1
10	340	4.9	200	4.1	100	3.0	4.3	3.1
11	360	4.9	200	4.2	100	3.0	4.4	3.0
12	370	4.9	200	4.2	100	3.2	3.7	3.0
13	390	4.8	200	4.2	100	3.2	3.5	3.0
14	380	4.8	210	4.1	100	3.1	3.3	2.9
15	370	4.9	220	4.0	100	3.0	3.8	3.0
16	340	5.0	220	3.8	110	2.8	4.1	3.0
17	300	5.3	230	3.6	110	2.5	4.3	3.1
18	280	5.8	240	3.2	110	1.9	4.7	3.1
19	250	6.0					4.4	3.2
20	240	5.7					4.3	3.3
21	240	4.6					3.5	3.2
22	250	3.7					3.0	3.2
23	270	3.0					3.0	3.1

Time: 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 2

Fairbanks, Alaska (64.9°N, 147.8°W)

March 1954

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	---	---					4.7	---
01	(340)	(2.2)					5.0	(3.0)
02	---	---					5.6	---
03	(310)	(2.4)					5.6	(3.0)
04	(350)	(2.0)					5.2	(2.9)
05	(360)	(2.0)					4.6	(2.9)
06	320	2.6					4.5	3.1
07	280	2.8	240	---	120	2.0	3.9	3.1
08	300	3.4	240	3.3	120	2.0	2.0	3.1
09	310	3.7	220	3.4	120	2.2	2.5	3.1
10	360	3.8	220	3.5	120	2.4		3.0
11	360	3.8	210	3.6	120	2.3		3.0
12	370	4.0	210	3.6	120	2.4		3.0
13	340	4.0	210	3.6	120	2.4		3.0
14	310	4.0	230	3.6	120	2.4		3.2
15	290	4.1	230	(3.5)	120	2.2		3.2
16	260	4.0	220	---	---	---		3.3
17	250	4.0	240	---	---	---		3.3
18	260	3.5						3.2
19	280	3.0					2.3	(3.1)
20	300	2.7					3.6	(3.0)
21	340	(2.2)					4.0	(3.1)
22	(350)	(2.3)					5.0	(3.0)
23	---	(2.1)					4.8	(2.9)

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 3

Anchorage, Alaska (61.2°N, 149.9°W)

March 1954

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	330	(2.0)					2.5	(2.95)
01	(360)	1.8					2.4	2.8
02	(350)	(2.0)					2.1	(2.9)
03	370	(2.3)					2.2	(2.8)
04	340	(1.8)					2.0	(2.8)
05	320	(2.2)					1.6	(2.9)
06	300	2.6	---	---	120	1.6		3.1
07	270	3.1	240	2.9	120	1.8		3.2
08	320	3.3	230	3.2	120	2.2		3.1
09	490	3.6	220	3.4	120	2.2		3.0
10	380	3.9	210	3.5	120	2.3		2.9
11	420	4.0	210	3.6	120	2.4		2.8
12	400	4.1	210	3.6	120	2.5		2.9
13	340	4.2	220	3.6	120	2.5		3.15
14	320	4.3	220	3.5	120	2.4		3.1
15	290	4.3	230	3.4	120	2.3		3.3
16	270	4.3	230	3.2	130	2.1		3.3
17	250	4.1	240	---	---	---		3.4
18	240	3.8						3.3
19	240	3.2						3.2
20	240	2.6						3.15
21	270	2.4						3.0
22	310	2.2						3.0
23	330	(2.2)						(3.0)

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 4

Adak, Alaska (51.9°N, 176.6°W)

March 1954

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	270	3.0						3.0
01	270	3.0						3.0
02	270	3.0						3.0
03	270	2.9						3.0
04	260	2.9						3.1
05	250	2.9						3.2
06	250	3.2	---	---	---	---		3.3
07	240	4.1	230	---	120	1.8		3.4
08	270	4.5	230	3.6	120	2.2		3.3
09	300	4.9	230	3.8	120	2.4	2.6	3.3
10	300	4.9	210	3.9	110	2.5	2.7	3.2
11	300	5.6	210	4.0	110	2.7		3.2
12	280	5.8	210	4.0	110	2.7		3.3
13	290	5.7	210	4.0	110	2.6		3.3
14	270	5.7	230	3.9	110	2.5		3.3
15	270	5.5	230	3.7	120	2.2		3.4
16	250	5.4	230	3.3	120	2.2		3.5
17	230	5.0	---	---	120	1.8		3.5
18	230	4.5			---	---	1.4	3.5
19	220	3.5						3.3
20	250	3.0						3.2
21	250	2.9						3.2
22	260	3.0						3.1
23	270	3.0						3.0

Time: 180.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 30 seconds.

Table 5

Graz, Austria (47.1°N, 15.5°E)

March 1954

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	2.9						
01	300	2.9						
02	300	2.9						
03	300	2.8						
04	290	2.6						
05	280	2.4						
06	280	2.6						
07	240	3.9	240	(3.1)				
08	230	4.3	200	3.5	110	(2.8)		
09	250	4.8	200	3.7	110	(2.9)		
10	260	5.0	200	3.9	110	(3.0)		
11	275	5.0	200	4.0	110	3.0	3.0	
12	260	5.4	200	4.0	110	2.9		
13	270	5.1	200	4.0	110	2.9		
14	260	5.1	200	4.0				
15	250	5.2	210	3.7				
16	240	5.1	220	3.5				
17	230	5.0	230	(3.1)				
18	220	4.9						
19	250	4.1						
20	250	3.8						
21	280	3.1						
22	280	2.9						
23	300	2.9						

Time: 15.0°E.

Sweep: 2.5 Mc to 12.0 Mc in 2 minutes.

Table 6

San Francisco, California (37.4°N, 122.2°W)

March 1954

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	(280)	(3.0)					2.2	3.0
01	(270)	3.0					2.4	3.0
02	(260)	3.1					2.2	3.1
03	260	(3.2)						(3.1)
04	(250)	3.0						3.2
05	260	3.0						3.1
06	250	(3.2)					(2.4)	(3.2)
07	260	4.3	250	---	(130)	(2.0)	(2.3)	3.4
08	280	4.9	230	(3.6)	120	(2.4)	3.4	3.3
09	310	5.2	220	(3.9)	120	(2.6)	3.6	3.2
10	330	5.3	210	(4.0)	110	(2.8)	3.8	3.0
11	320	5.8	210	(4.0)	120	(3.0)	3.9	3.0
12	300	6.3	220	(4.1)	(120)	(3.0)	3.7	3.1
13	320	6.0	230	(4.2)	(170)	(3.1)	3.6	3.1
14	300	5.8	230	(4.1)	120	(3.0)	3.8	3.2
15	300	5.5	230	(4.0)	120	(2.9)	3.6	3.2
16	280	5.2	240	(3.7)	120	(2.6)	2.7	3.3
17	250	5.2	---	---	(120)	(2.2)	2.9	3.4
18	240	4.7	---	---	---	---	1.9	3.4
19	230	3.8					2.4	3.3
20	240	3.3					3.2	3.2
21	(260)	(2.9)					2.5	(3.1)
22	(270)	3.1					2.4	3.0
23	(280)	3.0					2.5	3.0

Time: 120.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 7

White Sands, New Mexico (32.3°N, 106.5°W) March 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	300	3.3					3.0
01	280	3.3					3.0
02	270	3.3					3.0
03	260	3.3					3.1
04	270	3.2					3.05
05	270	3.0					3.1
06	260	3.5					3.1
07	250	4.5	240	---	120	1.9	3.3
08	270	5.2	220	3.7	120	2.3	3.7
09	300	5.3	210	4.0	120	2.6	3.9
10	310	5.5	200	4.1	120	3.0	3.8
11	310	6.0	200	4.2	120	3.0	4.0
12	310	6.4	200	4.2	120	3.2	3.9
13	310	6.7	210	4.3	120	3.2	3.4
14	300	6.4	220	4.2	120	3.1	3.9
15	300	5.8	220	4.0	120	2.9	3.9
16	280	5.8	220	3.8	120	2.6	3.7
17	260	5.6	230	3.2	120	2.2	3.3
18	230	5.3					2.9
19	230	4.4					2.4
20	250	3.5					2.6
21	270	3.2					3.05
22	280	3.2					3.0
23	300	3.2					3.0

Time: 105.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 8

Formosa, China (25.0°N, 121.5°E) March 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	290	3.4					2.0
01	260	3.5					2.1
02	250	3.7					1.8
03	240	3.4					3.4
04	240	2.6					3.2
05	260	2.1					3.0
06	260	2.8					3.1
07	240	5.3			120	2.0	3.5
08	250	6.1	230	3.9	120	2.6	3.2
09	280	6.9	240	4.2	120	2.9	3.8
10	300	7.6	230	4.3	120	3.1	3.7
11	310	9.2	230	4.4	120	3.2	3.8
12	290	10.8	220	4.4	170	(3.3)	4.4
13	280	11.2	220	4.4	120	(3.3)	4.2
14	280	11.6	230	4.3	120	3.2	4.2
15	270	11.4	230	4.1	110	3.1	4.1
16	240	10.2	220	3.8	120	2.8	3.7
17	240	8.6	220	3.4	120	2.4	3.7
18	230	6.8			---	---	3.2
19	230	5.6					3.3
20	240	4.8					2.2
21	270	4.0					2.0
22	280	3.5					1.9
23	300	3.8					2.9

Time: 120.0°E.

Sweep: 1.1 Mc to 19.5 Mc in 15 minutes, manual operation.

Table 9

Maui, Hawaii (20.8°N, 156.5°W) March 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	270	3.2					1.3
01	270	3.3					1.1
02	250	3.2					1.2
03	230	2.7					1.6
04	250	2.0					3.35
05	270	1.9					3.3
06	300	1.9					3.0
07	250	4.5	240	---	140	1.7	2.4
08	(270)	5.7	240	---	120	2.4	3.3
09	320	6.6	230	4.2	120	2.8	4.2
10	340	8.0	210	4.4	120	3.0	4.6
11	360	8.9	220	4.4	120	3.2	4.5
12	340	10.0	210	4.5	120	3.3	4.8
13	320	11.2	210	4.5	120	3.3	4.5
14	300	11.6	210	4.4	120	3.2	4.6
15	280	11.2	230	4.3	120	3.0	4.0
16	270	10.7	230	4.1	120	2.8	3.8
17	250	9.0	240	---	120	2.4	3.7
18	240	6.6			130	1.6	3.4
19	240	5.8					3.5
20	240	4.5					2.2
21	250	3.9					1.9
22	280	3.3					2.0
23	290	3.5					1.5

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 10

Puerto Rico, W. I. (18.5°N, 67.2°W) March 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	270	3.7					3.1
01	260	3.8					3.1
02	240	3.7					3.2
03	230	3.7					3.3
04	230	3.2					1.9
05	230	2.7					3.3
06	250	2.5					3.1
07	220	4.2	---	---	---	---	3.55
08	240	5.2	230	---	110	2.3	2.8
09	270	5.8	220	4.0	110	2.7	3.2
10	290	6.5	220	4.3	110	3.0	2.7
11	300	7.0	220	4.3	110	3.2	3.5
12	290	7.7	210	4.4	110	3.3	3.4
13	280	8.0	210	4.4	110	3.3	3.2
14	280	8.3	210	4.3	110	3.2	2.4
15	270	8.4	220	4.2	110	3.1	2.9
16	250	7.7	220	4.0	110	2.9	4.1
17	250	6.7	220	---	110	2.5	3.6
18	230	6.5	230	---	110	---	2.9
19	210	5.9					2.4
20	220	4.6					3.3
21	250	3.9					3.1
22	270	3.7					3.0
23	280	3.6					3.0

Time: 60.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 11

Guam I. (13.6°N, 144.3°E) March 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	260	4.3					2.5
01	260	4.2					1.6
02	260	3.6					1.6
03	260	3.0					3.2
04	< 260	2.4					3.3
05	250	1.9					1.8
06	240	1.7					1.8
07	240	5.1	240	---	120	(1.5)	2.8
08	260	6.2	230	---	120	2.5	3.8
09	300	7.2	220	4.1	110	2.9	4.8
10	320	7.9	210	4.2	110	3.1	4.7
11	350	8.4	200	4.3	110	(3.2)	4.7
12	350	8.2	200	4.3	110	3.2	4.3
13	360	8.0	200	4.3	110	3.2	4.4
14	340	8.5	200	4.2	110	3.2	4.5
15	320	9.3	220	4.2	110	3.0	4.4
16	300	9.5	230	4.1	110	(2.7)	3.8
17	270	9.9	230	---	120	2.4	3.9
18	250	9.6	250	---	---	---	3.4
19	240	9.0					2.6
20	240	8.0					2.4
21	230	6.6					2.7
22	240	5.7					2.8
23	260	4.6					2.6

Time: 150.0°E.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 12

Kiruna, Sweden (67.8°N, 20.3°E) February 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	---	---					2.6
01	---	---					---
02	---	---					---
03	(350)	(2.0)					(3.0)
04	---	---					---
05	---	---					---
06	---	---					---
07	290	2.0					3.3
08	250	3.1	---	---	---	---	3.5
09	250	3.4	---	---	---	---	3.7
10	250	3.7	---	---	---	---	3.6
11	250	3.9	220	2.8	---	---	3.6
12	250	4.3	210	---	---	---	3.6
13	250	4.2	---	---	---	---	3.6
14	250	4.1	---	---	---	---	3.6
15	230	3.8	---	---	---	---	3.6
16	240	3.2	---	---	---	---	3.6
17	(250)	(2.5)					2.0
18	---	---					3.1
19	---	---					3.4
20	---	---					2.8
21	---	---					4.0
22	---	---					3.1
23	---	---					3.3

Time: 15.0°E.

Sweep: 0.8 Mc to 15.0 Mc in 30 seconds.

Table 13

Fairbanks, Alaska (64.9°N, 147.8°W) February 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	---	---	---	---	---	---	5.0
01	---	---	---	---	---	---	5.1
02	---	---	---	---	---	---	5.5
03	---	---	---	---	---	---	5.7
04	---	---	---	---	---	---	5.4
05	---	---	---	---	---	---	4.5
06	---	---	---	---	---	---	4.5
07	---	---	---	---	---	---	4.5
08	(1.9)	---	---	---	---	---	3.4
09	2.6	---	---	---	---	---	2.5
10	3.4	---	---	---	---	---	2.2
11	3.8	---	---	---	---	---	1.9
12	4.0	---	---	---	---	---	2.0
13	4.3	---	---	---	---	---	2.8
14	4.4	---	---	---	---	---	---
15	4.6	---	---	---	---	---	2.0
16	4.3	---	---	---	---	---	1.8
17	4.0	---	---	---	---	---	1.8
18	3.6	---	---	---	---	---	(4.0)
19	2.8	---	---	---	---	---	3.6
20	(1.9)	---	---	---	---	---	3.4
21	(1.8)	---	---	---	---	---	5.0
22	(2.0)	---	---	---	---	---	3.8
23	(2.0)	---	---	---	---	---	3.8
23	(2.2)	---	---	---	---	---	4.4

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 14

Adak, Alaska (51.9°N, 176.6°W) February 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	---	---	---	---	---	---	2.7
01	---	---	---	---	---	---	2.7
02	---	---	---	---	---	---	2.7
03	---	---	---	---	---	---	2.7
04	---	---	---	---	---	---	2.6
05	---	---	---	---	---	---	2.5
06	---	---	---	---	---	---	2.7
07	---	---	---	---	---	---	3.2
08	---	---	---	---	---	---	4.1
09	---	---	---	---	---	---	4.9
10	---	---	---	---	---	---	3.1
11	---	---	---	---	---	---	5.4
12	---	---	---	---	---	---	3.6
13	---	---	---	---	---	---	3.8
14	---	---	---	---	---	---	3.8
15	---	---	---	---	---	---	5.5
16	---	---	---	---	---	---	5.6
17	---	---	---	---	---	---	5.2
18	---	---	---	---	---	---	4.4
19	---	---	---	---	---	---	3.1
20	---	---	---	---	---	---	2.5
21	---	---	---	---	---	---	2.3
22	---	---	---	---	---	---	2.4
23	---	---	---	---	---	---	2.5
23	---	---	---	---	---	---	2.7

Time: 180.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 30 seconds.

Table 15

Guam I. (13.6°N, 144.9°E) February 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	240	4.1	---	---	---	---	3.2
01	250	4.0	---	---	---	---	3.2
02	250	3.6	---	---	---	---	3.2
03	240	3.1	---	---	---	---	3.4
04	260	2.6	---	---	---	---	3.4
05	260	2.2	---	---	---	---	1.7
06	280	1.3	---	---	---	---	1.7
07	250	4.1	240	---	---	---	(3.4)
08	260	6.0	230	---	---	---	3.5
09	290	7.3	220	---	---	---	3.4
10	320	8.0	210	---	---	---	2.7
11	340	8.0	200	---	---	---	120
12	350	7.7	200	---	---	---	4.0
13	350	7.4	200	---	---	---	110
14	340	8.0	200	---	---	---	110
15	330	8.4	220	---	---	---	110
16	300	8.8	220	---	---	---	110
17	280	9.0	240	---	---	---	110
18	250	8.8	---	---	---	---	110
19	240	8.6	---	---	---	---	120
20	230	7.6	---	---	---	---	2.4
21	230	6.2	---	---	---	---	2.8
22	240	5.1	---	---	---	---	2.6
23	240	4.4	---	---	---	---	2.6
23	240	4.4	---	---	---	---	2.4

Time: 150.0°E.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 16

Leopoldville, Belgian Congo (4.3°S, 15.3°E) February 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M2000)F2
00	230	3.4	---	---	---	---	2.3
01	260	3.2	---	---	---	---	2.3
02	250	3.1	---	---	---	---	2.4
03	250	2.9	---	---	---	---	2.4
04	250	2.5	---	---	---	---	1.4
05	240	3.0	---	---	---	---	1.6
06	255	5.2	230	---	---	---	1.8
07	270	6.2	220	---	---	---	2.6
08	310	6.5	220	---	---	---	2.6
09	360	7.3	215	---	---	---	2.6
10	365	8.2	210	---	---	---	3.0
11	370	9.0	200	---	---	---	3.0
12	380	9.2	210	---	---	---	3.2
13	350	10.0	200	---	---	---	3.4
14	350	10.0	210	---	---	---	3.5
15	335	10.4	220	---	---	---	3.5
16	310	10.3	240	---	---	---	2.8
17	270	10.1	250	---	---	---	2.3
18	245	< 9.2	---	---	---	---	2.3
19	245	8.6	---	---	---	---	2.2
20	245	8.2	---	---	---	---	2.4
21	230	8.1	---	---	---	---	2.4
22	215	8.1	---	---	---	---	2.5
23	200	5.6	---	---	---	---	2.8
23	200	5.6	---	---	---	---	2.7

Time: 0.0°.

Sweep: 1.0 Mc to 16.0 Mc in 7 seconds.

Table 17

Johannesburg, Union of S. Africa (26.2°S, 28.1°E) February 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	260	3.4	---	---	---	---	3.0
01	250	3.4	---	---	---	---	3.1
02	240	3.3	---	---	---	---	1.8
03	240	3.1	---	---	---	---	3.2
04	250	2.8	---	---	---	---	2.7
05	250	2.7	---	---	---	---	3.1
06	250	3.6	---	---	---	---	2.2
07	260	4.7	230	---	---	---	3.4
08	320	5.1	220	---	---	---	3.4
09	330	5.6	210	---	---	---	3.6
10	320	6.2	200	---	---	---	3.1
11	330	6.3	200	---	---	---	4.3
12	340	6.4	200	---	---	---	4.3
13	330	6.6	200	---	---	---	4.4
14	320	6.4	210	---	---	---	4.4
15	320	6.8	220	---	---	---	4.3
16	300	6.6	220	---	---	---	4.1
17	280	6.2	220	---	---	---	4.0
18	260	6.1	230	---	---	---	3.7
19	240	6.0	---	---	---	---	3.7
20	240	5.6	---	---	---	---	2.2
21	240	4.7	---	---	---	---	2.0
22	240	3.8	---	---	---	---	3.2
23	260	3.5	---	---	---	---	3.1
23	260	3.5	---	---	---	---	3.0

Time: 30.0°E.

Sweep: 1.0 Mc to 15.0 Mc in 7 seconds.

Table 18

Capetown, Union of S. Africa (34.2°S, 18.3°E) February 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	250	3.2	---	---	---	---	3.1
01	< 260	3.2	---	---	---	---	3.0
02	260	3.2	---	---	---	---	3.0
03	< 260	3.2	---	---	---	---	2.2
04	250	3.2	---	---	---	---	2.0
05	< 250	3.1	---	---	---	---	3.1
06	250	3.0	---	---	---	---	3.1
07	250	4.0	240	---	---	---	3.0
08	280	4.7	240	---	---	---	3.6
09	320	5.2	230	---	---	---	4.0
10	340	5.4	220	---	---	---	4.1
11	340	5.9	220	---	---	---	4.2
12	340	6.0	200	---	---	---	4.2
13	340	6.3	200	---	---	---	4.3
14	340	6.2	220	---	---	---	4.2
15	330	6.1	210	---	---	---	4.2
16	310	6.2	220	---	---	---	4.1
17	290	5.8	220	---	---	---	3.9
18	290	5.6	220	---	---	---	3.5
19	250	5.4	240	---	---	---	2.9
20	240	5.0	---	---	---	---	2.2
21	240	4.8	---	---	---	---	2.1
22	240	4.0	---	---	---	---	1.8
23	250	3.6	---	---	---	---	1.7
23	250	3.6	---	---	---	---	3.2

Time: 30.0°E.

Sweep: 1.0 Mc to 15.0 Mc in 7 seconds.

Table 19

Resolute Bay, Canada (74.7°N, 94.9°W)								January 1954	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	260	2.3						3.2	
01	250	2.3						3.2	
02	250	2.0						3.2	
03	270	2.0					2.7	3.2	
04	270	2.1					3.8	3.2	
05	270	2.4					3.5	3.2	
06	280	2.1					3.4	3.2	
07	250	2.5					1.8	3.2	
08	260	2.6					3.1	3.2	
09	250	3.0					3.7	3.2	
10	250	3.0					3.2	3.2	
11	250	3.0			120	1.0	3.4	3.2	
12	240	3.4			120	1.1	1.9	3.2	
13	240	3.1			120	1.1		3.2	
14	230	3.2						3.2	
15	250	3.1						3.2	
16	240	3.0						3.2	
17	240	3.1						3.2	
18	240	3.0						3.2	
19	250	3.0						3.2	
20	250	2.8						3.2	
21	260	2.4						3.2	
22	260	2.7						3.2	
23	250	2.6						3.2	

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 20

Fairbanks, Alaska (64.9°N, 147.8°W)								January 1954	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00		(2.1)					4.5	(3.0)	
01		(2.1)					4.1	(3.0)	
02		(2.3)					4.7	(3.0)	
03		(2.1)					4.5	(2.8)	
04		(2.0)					4.5	(2.8)	
05		(1.9)					4.0	(2.8)	
06		(1.8)					4.0	(2.9)	
07		(1.8)					4.5	(2.8)	
08		(1.8)					4.4	(3.0)	
09		2.8					2.5	3.3	
10		3.7					2.5	3.5	
11		4.2					2.3	3.5	
12		4.5					1.9	3.5	
13		4.7					2.3	3.5	
14		4.6					3.6	3.5	
15		4.2					2.4	3.5	
16		3.4					3.8	3.4	
17		2.6					4.0	3.4	
18		2.1					3.8	3.4	
19		(1.6)					4.6	(3.2)	
20		---					4.0	---	
21		(2.0)					3.7	(3.2)	
22		(2.5)					4.0	(3.3)	
23		(2.5)					4.2	(3.1)	

Time: 150.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 21

Baker Lake, Canada (64.3°N, 96.0°W)								January 1954	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	250	2.0			---	E	7.0	3.2	
01	260	2.0			---	E	6.0	3.2	
02	250	1.8			---	E	5.2	3.2	
03	260	1.8			---	E	4.8	3.2	
04	260	2.3			---	1.1	4.8	(3.3)	
05	250	2.4			---	1.3	4.0	(3.2)	
06	250	2.6			130	1.6	3.8	(3.2)	
07	250	2.8			120	2.1	4.1	(3.2)	
08	260	3.2			110	2.3	4.4	(3.2)	
09	270	3.6			110	2.4	4.8	3.2	
10	270	4.0			110	3.0	3.4	3.2	
11	260	4.1			100	2.7	3.3	3.3	
12	260	4.3			110	2.8	2.3	3.3	
13	250	4.8			110	2.6		3.3	
14	250	5.0			110	2.3		3.3	
15	240	4.5			120	2.2	2.6	3.3	
16	250	3.8			120	2.2	2.3	3.2	
17	280	3.3			120	2.2	2.6	(3.1)	
18	260	3.4			120	2.1	4.0	3.1	
19	250	3.2			120	2.1	5.0	(3.0)	
20	250	3.0			130	1.7	6.0	3.0	
21	240	2.6			---	E	6.4	(3.1)	
22	240	2.7			---	E	7.0	3.0	
23	250	2.4			---	E	7.5	3.2	

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 22

Churchill, Canada (58.8°N, 94.2°W)								January 1954	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	---	---			---	---	7.0	---	
01	---	---			---	---	6.0	---	
02	---	---			---	---	5.0	---	
03	---	---			---	---	5.0	---	
04	---	---			---	---	4.5	---	
05	---	---			---	---	4.6	---	
06	---	---			---	---	5.0	---	
07	---	---			---	---	5.4	---	
08	(300)	(3.1)			---	---	5.1	---	
09	260	3.5			110	2.0	4.6	3.4	
10	260	4.1			120	2.1	3.2	3.4	
11	250	4.4			110	2.4		3.4	
12	260	4.8	220		120	2.4		3.4	
13	260	5.0	220		120	2.2		3.3	
14	250	5.2	220		120	2.1		3.4	
15	240	5.0	---		120	2.0		3.4	
16	230	4.8			120	1.9		3.4	
17	250	4.0			---	---	3.5	3.3	
18	270	3.2			120	---	3.8	3.3	
19	290	2.7			---	---	3.7	3.2	
20	320	2.8			---	---	4.1	(3.1)	
21	(320)	(2.8)			---	---	4.8	---	
22	---	---			---	---	4.6	---	
23	---	---			---	---	5.8	---	

Time: 90.0°W.

Sweep: 0.6 Mc to 10.0 Mc in 16 seconds.

Table 23

Fort Chino, Canada (58.1°N, 68.3°W)								January 1954	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	---	<2.5			110	2.5	4.2		
01	---	2.3			110	2.7	4.4		
02	---	<3.0			100	3.0	4.2		
03	---	---			100	3.4	4.3		
04	---	---			100	3.1	4.3	---	
05	---	---			100	3.2	4.0		
06	---	---			100	2.7	4.0		
07	---	<2.0			100	2.3	3.5		
08	240	3.3			100	1.8		(3.5)	
09	230	4.0	210		100	2.0		3.5	
10	230	4.4	210		100	2.2		3.5	
11	240	4.9	210		110	2.2		3.5	
12	240	5.2	200		110	2.3		3.4	
13	240	5.2	220	(3.1)	110	2.2		3.4	
14	230	5.2	210		120	2.0		3.4	
15	240	5.0			110	2.1		(3.4)	
16	240	4.2			100	2.1		---	
17	260	3.1			100	3.0			
18	270	2.8			100	3.0	2.2		
19	280	<2.6			100	2.9	3.2		
20	260	3.0			100	2.4	5.2		
21	---	<2.8			---	---	6.1		
22	---	---			100	3.6	4.3		
23	---	---			100	2.9	5.1		

Time 75.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 24

Prince Rupert, Canada (54.3°N, 130.3°W)								January 1954	
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2	
00	300	1.5					2.4		
01	290	1.4					2.2		
02	300	1.5					> 3.0		
03	300	1.5					3.0	---	
04	290	1.5					4.0		
05	300	1.6					3.1		
06	300	1.8					4.0		
07	290	1.5					3.4		
08	270	1.9					3.0	---	
09	240	3.4			110		3.0	3.4	
10	230	4.4	210		110	1.9	2.1	3.5	
11	230	4.9	220		110	2.1		3.5	
12	240	5.2	220	3.3	110	2.2		3.5	
13	240	5.3	210	3.3	110	2.3		3.5	
14	240	5.7	220		120	2.2		3.5	
15	230	5.3	220		120	2.0		3.5	
16	210	4.7			120	1.8		3.5	
17	210	4.0				E		3.4	
18	210	3.1						3.4	
19	250	2.1						---	
20	270	1.7						---	
21	(260)	1.5					1.2	---	
22	---	1.6					2.1		
23	---	1.6							

Time: 120.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 15 seconds.

Table 25

Winnipeg, Canada (49.9°N, 97.4°W)							
January 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	380	(2.3)					---
01	350	(2.3)					---
02	(300)	(2.5)					3.5
03	350	(2.5)					3.5
04	(300)	(2.4)					3.8
05	340	(2.2)					3.4
06	---	(2.3)					3.9
07	(320)	(2.4)					3.9
08	260	2.4					3.4
09	230	3.9	210	---	120	1.9	3.2
10	240	4.6	220	---	120	2.1	3.4
11	250	5.0	220	---	120	2.3	3.4
12	260	5.3	220	3.6	120	2.5	3.4
13	260	5.6	220	3.5	120	2.5	3.4
14	250	5.8	220	---	120	2.4	3.4
15	240	5.6	230	---	120	2.2	3.4
16	230	5.2	220	---	130	1.9	3.5
17	220	4.4					3.4
18	230	3.5					3.3
19	250	2.5					3.2
20	300	1.9					3.0
21	(350)	1.8					---
22	(480)	(2.2)					---
23	(400)	(2.4)					---

Time: 90.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 16 seconds.

Table 26

Ottawa, Canada (45.4°N, 75.9°W)							
January 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	360	1.6					---
01	(340)	1.6					---
02	360	1.8					(3.1)
03	350	1.9					(3.1)
04	310	2.0					(3.2)
05	290	2.0					2.2
06	300	2.0					3.0
07	290	2.2					2.5
08	230	3.7	---	---	160	1.9	3.1
09	230	4.6	220	2.6	120	2.1	3.2
10	250	5.0	210	3.4	120	2.4	3.5
11	260	5.4	210	3.7	120	2.6	3.4
12	250	5.4	220	3.8	120	2.7	3.5
13	260	5.5	220	3.8	120	2.7	3.4
14	260	5.7	230	3.5	120	2.5	3.5
15	250	5.3	230	3.2	120	2.3	3.5
16	230	5.0	230	---	130	1.9	3.4
17	220	4.5					3.4
18	240	3.6					3.3
19	250	3.0					3.3
20	270	2.3					3.3
21	300	2.0					3.2
22	360	1.8					(3.2)
23	360	1.7					---

Time: 75.0°W.

Sweep: 1.0 Mc to 10.0 Mc in 15 seconds.

Table 27

Johannesburg, Union of S. Africa (26.2°S, 28.1°E)							
January 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	250	3.5					2.7
01	250	3.3					3.1
02	250	3.0					2.6
03	250	2.9					2.6
04	250	2.7					2.4
05	250	2.7					2.0
06	240	4.0	250	---	130	1.8	2.0
07	290	4.7	230	3.6	110	2.3	2.4
08	330	5.2	220	4.0	110	2.8	3.1
09	340	5.7	210	4.1	110	3.1	3.8
10	360	5.8	200	4.3	110	3.2	3.9
11	350	6.4	200	4.4	110	3.3	4.1
12	340	6.8	200	4.4	110	3.4	4.2
13	340	7.2	200	4.4	110	3.4	4.2
14	320	7.1	200	4.3	110	3.3	4.0
15	310	7.1	210	4.2	110	3.2	3.8
16	290	6.9	210	4.0	110	2.9	3.6
17	270	6.2	210	3.7	110	2.6	3.6
18	260	5.4	220	3.2	120	2.1	3.1
19	240	5.2					3.0
20	240	5.3					2.2
21	240	4.7					2.5
22	240	4.1					2.4
23	260	3.6					2.4

Time: 30.0°E.

Sweep: 1.0 Mc to 15.0 Mc in 7 seconds.

Table 28

Capetown, Union of S. Africa (34.2°S, 18.3°E)							
January 1954							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
000	250	3.3					2.6
01	<260	3.2					3.0
02	260	3.2					2.6
03	260	3.2					3.0
04	260	3.0					3.0
05	250	3.0					2.6
06	240	3.6	250	---	140	1.6	3.0
07	280	4.4	240	3.4	120	2.1	2.1
08	340	5.0	230	3.8	110	2.6	3.2
09	350	5.2	220	4.0	110	2.9	3.0
10	360	5.5	210	4.2	110	3.1	3.0
11	360	5.9	210	4.3	110	3.2	4.2
12	350	6.4	200	4.3	110	3.3	2.9
13	340	6.5	200	4.3	110	3.3	4.4
14	330	6.6	200	4.3	110	3.3	4.8
15	320	6.4	200	4.2	110	3.2	4.3
16	320	6.0	210	4.1	110	3.1	4.0
17	300	5.6	220	4.0	110	2.9	3.0
18	290	5.5	210	3.7	110	2.5	3.8
19	260	5.2	220	3.0	120	2.0	3.3
20	230	5.2					3.0
21	230	5.0					2.5
22	240	4.0					2.3
23	240	3.6					2.6

Time: 30.0°E.

Sweep: 1.0 Mc to 15.0 Mc in 7 seconds.

Table 29

Baker Lake, Canada (64.3°N, 96.0°W)							
December 1953							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	250	2.2					8.0
01	260	2.4					7.0
02	270	2.4					6.0
03	260	2.4					5.0
04	260	2.5					1.4
05	250	2.6					1.6
06	250	2.7					1.8
07	250	3.0					1.8
08	260	3.0					2.3
09	250	3.4					2.6
10	280	3.6					2.7
11	260	4.0					2.6
12	260	4.2					2.3
13	250	4.8					2.2
14	250	5.0					2.0
15	240	4.2					2.0
16	260	3.5					2.3
17	260	3.3					2.4
18	260	3.4					2.3
19	250	3.3					2.0
20	250	2.9					2.0
21	250	2.8					5.8
22	250	2.5					6.5
23	260	2.3					7.2

Time: 90.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 15 seconds.

Table 30

Reykjavik, Iceland (64.1°N, 21.8°W)							
December 1953							
Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs (M3000)F2
00	---	---					4.6
01	---	---					4.2
02	---	(2.9)					4.0
03	320	(2.4)					4.1
04	330	(2.4)					1.8
05	335	2.3					(3.1)
06	---	---					3.0
07	---	---					(3.9)
08	---	---					(3.7)
09	280	2.5					3.2
10	250	3.0					3.4
11	235	3.6					3.4
12	240	4.0					3.4
13	240	4.2					3.4
14	240	4.1					3.4
15	240	3.7					3.2
16	260	3.0					3.0
17	260	(2.6)					3.6
18	---	(2.4)					3.9
19	---	---					4.3
20	---	---					4.0
21	---	---					4.4
22	---	---					5.1
23	---	---					4.7

Time: 15.0°W.

Sweep: 1.0 Mc to 25.0 Mc in 18 seconds.

Table 31

Wakkanai, Japan (45.4°N, 141.7°E) December 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	270	3.2					2.4	3.0
01	280	3.3					2.4	3.0
02	270	3.1					2.4	3.1
03	250	3.0					2.2	3.1
04	250	2.9					1.6	3.2
05	230	2.9					2.4	3.4
06	250	2.6					2.6	3.4
07	220	3.5					2.7	3.6
08	220	4.7				2.0	2.7	3.7
09	230	5.3	240	3.4	120	2.2	3.2	3.6
10	230	6.0	230	3.6	120	2.5	3.3	3.6
11	230	5.8	230	3.6	120	2.6	2.6	3.6
12	230	5.6	230	3.6	120	2.6	2.6	3.6
13	230	5.5	230	3.5	120	2.5	2.6	3.6
14	230	5.4	220	---	120	2.3	2.5	3.6
15	220	5.0			120	---	2.0	3.6
16	210	4.2						3.5
17	230	3.2						3.3
18	250	2.8					2.2	3.3
19	250	2.8					2.8	3.3
20	250	2.7					2.8	3.2
21	260	2.8					3.0	3.1
22	260	3.0					2.7	3.1
23	280	3.1					2.7	3.1

Time: 135.0°E.

Sweep: 1.0 Mc to 15.5 Mc in 2 minutes.

Table 32

Akita, Japan (39.7°N, 140.1°E) December 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	290	3.0					3.0	2.9
01	280	3.1					2.7	2.9
02	280	3.1					2.6	2.8
03	250	3.0					2.6	3.0
04	250	2.8					2.4	3.0
05	240	2.8					2.6	3.2
06	240	2.4					2.4	3.2
07	220	3.8					2.3	3.4
08	220	4.7	220	---	120	2.0	3.2	3.6
09	240	5.3	220	3.3	110	2.4	4.2	3.5
10	250	6.0	240	3.7	110	2.6	4.2	3.4
11	240	6.6	220	3.7	110	2.7	3.5	3.6
12	240	5.7	220	3.7	110	2.7	3.5	3.5
13	240	5.5	230	3.6	120	2.6	3.3	3.5
14	240	5.5	240	3.5	120	2.5	3.5	3.5
15	230	5.4	220	---	120	2.1	3.1	3.6
16	220	4.7					3.0	3.5
17	210	3.2					2.8	3.4
18	240	2.8					2.8	3.3
19	240	2.8					3.0	3.4
20	250	2.7					3.2	3.2
21	250	2.8					3.1	3.1
22	260	3.0					3.2	3.0
23	280	3.0					3.1	2.8

Time: 135.0°E.

Sweep: 0.85 Mc to 22.0 Mc in 2 minutes.

Table 33

Tokyo, Japan (35.7°N, 139.5°E) December 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	290	2.8					2.8	3.0
01	280	2.9					2.8	2.9
02	270	3.0					2.8	3.0
03	250	2.9					2.7	3.1
04	240	2.9					2.7	3.1
05	240	2.6					2.6	3.1
06	250	2.4					2.5	3.1
07	220	4.3			160	1.7	2.7	3.5
08	220	5.2	220	---	120	2.2	2.9	3.5
09	240	5.2	230	3.6	120	2.5	3.3	3.4
10	250	5.7	220	3.9	120	2.7	3.8	3.4
11	250	6.4	230	4.0	120	2.8	4.5	3.5
12	250	5.8	220	4.0	110	2.8	3.3	3.4
13	250	6.1	230	3.8	110	2.8	3.5	3.4
14	250	5.6	230	3.6	120	2.6	3.4	3.4
15	230	5.5	230	3.0	120	2.3	3.0	3.5
16	220	5.0	---	---	130	1.7	3.0	3.5
17	220	3.7					2.8	3.4
18	240	2.8					3.0	3.1
19	240	3.0					3.2	3.2
20	240	2.3					2.7	3.2
21	250	2.7					3.2	3.1
22	250	2.7					2.8	3.0
23	280	2.7					2.8	2.9

Time: 135.0°E.

Sweep: 1.0 Mc to 17.2 Mc in 2 minutes.

Table 34

Yamagawa, Japan (31.2°N, 130.6°E) December 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	320	2.5					2.0	2.9
01	310	2.6					1.8	2.9
02	300	2.7					2.0	3.0
03	290	2.7					1.8	3.1
04	260	2.7					2.0	3.1
05	270	2.5					2.0	3.2
06	290	2.4					2.0	3.0
07	250	3.2					2.2	3.2
08	240	4.9			130	2.0	2.2	3.6
09	250	5.1	240	---	120	2.3	2.0	3.5
10	260	5.8	240	3.9	110	2.7	3.4	3.4
11	260	6.0	250	4.0	120	2.8	3.4	3.3
12	270	6.6	220	4.0	110	2.9	4.0	3.4
13	270	6.4	220	4.0	110	2.8	3.6	3.4
14	280	6.4	250	3.9	---	2.8	3.6	3.4
15	260	6.0	250	3.7	110	2.5	3.6	3.5
16	250	5.8	230	3.1	120	2.2	3.4	3.5
17	240	4.9					3.4	3.6
18	240	3.5					2.6	3.4
19	260	2.9					2.5	3.1
20	260	2.8					2.6	3.1
21	260	2.6					2.5	3.1
22	270	2.6					2.4	3.1
23	300	2.5					2.1	3.1

Time: 135.0°E.

Sweep: 0.8 Mc to 20.0 Mc in 15 minutes, manual operation.

Table 35

Calcutta, India (22.6°N, 88.4°E) December 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	210	4.6						3.3
01	(210)	(4.2)						
02	(240)	(4.1)						
03	(225)	(3.6)						3.2
04	(240)	(3.2)						
05	(210)	(2.4)						
06	(210)	3.2						3.05
07	(220)	(4.8)						
08	210	6.5				2.4		
09	210	7.7				2.7		3.05
10	210	8.8				3.0		
11	210	10.0				3.2		
12	(210)	(9.6)				3.1		3.05
13	(210)	(9.4)				3.2		
14	(210)	(9.6)				3.0		
15	(230)	9.6				3.0		3.0
16	(240)	(9.0)				---		
17	(210)	(8.8)						
18	(210)	(9.6)						3.25
19	(210)	(5.2)						
20	(210)	(4.6)						
21	(240)	(4.5)						3.1
22	(210)	(4.1)						
23	(210)	(4.0)						

Time: 90.0°E.

Sweep: 0.5 Mc to 18.0 Mc in 10 minutes, semi-automatic operation.

Table 36

Barotonga I. (21.3°S, 159.8°W) December 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	270	6.2					3.9	3.0
01	270	5.5					3.2	3.0
02	280	4.8					3.0	3.15
03	300	3.9					3.0	3.0
04	290	4.1					2.9	2.9
05	300	3.9					2.5	2.95
06	250	4.7				1.9	3.1	3.2
07	260	6.0	230	3.8	110	2.4	3.7	3.3
08	290	6.5	230	4.2	105	2.8	4.4	3.1
09	300	7.0	200	4.3	105	3.1	4.5	3.2
10	320	7.8	200	4.3	105	3.3	4.7	3.1
11	330	8.7	200	4.4	105	3.4	4.5	2.9
12	340	9.4	200	4.4	105	3.4	5.4	2.85
13	320	10.3	200	4.4	105	3.4	4.7	3.0
14	310	10.3	200	4.3	105	3.4	4.5	3.0
15	310	9.7	---	---	105	3.2	4.7	3.0
16	290	9.3	---	---	110	3.0	4.2	3.1
17	280	9.1	240	3.8	110	2.6	5.2	3.0
18	260	8.3			125	---	4.1	3.1
19	260	7.6					4.0	3.0
20	290	7.1					3.5	2.9
21	300	6.4					3.5	2.9
22	300	6.5					3.0	2.9
23	290	6.0					3.2	2.9

Time: 157.5°W.

Sweep: 2.0 Mc to 16.0 Mc, manual operation.

Table 37*

Inverness, Scotland (57.4°N, 4.2°W)

November 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	345	(1.8)						(2.8)
01	330	1.6						2.8
02	315	1.5						2.9
03	330	1.5					2.3	2.8
04	320	(1.5)					2.5	2.9
05	305	(1.5)					2.4	(2.9)
06	(270)	(1.5)					2.4	(3.1)
07	290	(1.8)					2.4	3.0
08	240	2.7					2.4	3.4
09	225	3.8	(220)		(125)	1.8	2.4	3.6
10	240	4.6	215	(3.1)	(125)	2.0	2.4	3.5
11	240	4.9	215	3.2	(114)	2.2	2.4	3.5
12	240	5.1	215	3.2	120	2.2	2.4	3.5
13	245	5.2	220	3.2	120	2.1	2.4	3.5
14	240	5.2	225	(3.0)	135	2.0	2.0	3.5
15	235	5.0			150	1.8	2.1	3.4
16	220	4.3					2.4	3.4
17	230	3.5					2.6	3.2
18	240	2.5						3.3
19	265	1.9						3.3
20	290	(1.7)						3.1
21	320	(1.7)						3.0
22	350	(1.7)						(2.9)
23	345	(1.6)						

Time: 0.0°.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 38*

Slough, England (51.5°N, 0.6°W)

November 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	275	2.9						2.4
01	270	2.9						2.4
02	265	2.8						2.5
03	265	2.4						2.6
04	270	2.2						2.6
05	255	2.0						2.9
06	265	2.0						2.5
07	245	2.5						2.5
08	225	4.0			130	1.7		3.2
09	230	4.8	220	3.0	125	2.0		3.2
10	245	5.0	215	3.4	125	2.3		3.6
11	250	5.8	215	3.6	125	2.5		3.5
12	250	5.8	220	3.6	125	2.5		3.5
13	240	5.6	215	3.4	125	2.4		3.6
14	245	5.3	225	3.1	125	2.2		3.5
15	235	5.3			130	2.0		3.4
16	225	4.6						3.0
17	230	3.8						3.0
18	240	3.2						2.6
19	260	2.6						2.4
20	280	2.4						2.4
21	285	2.4						2.4
22	290	2.6						2.4
23	285	2.8						2.2

Time: 0.0°.

Sweep: 0.55 Mc to 16.5 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 39

Delhi, India (28.6°N, 77.1°E)

November 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	280	2.6						3.2
01	280	2.7						3.3
02	--	2.5						3.2
03	--	--						---
04	240	3.0						3.6
05	250	3.0						3.5
06	240	3.3						3.6
07	220	5.6						3.8
08	240	6.8						3.6
09	240	7.0						3.6
10	240	7.2						3.6
11	240	7.5						3.6
12	240	7.0						3.6
13	260	7.0						3.5
14	240	7.5						3.6
15	240	7.2						3.6
16	220	6.9						3.8
17	220	6.0						3.8
18	220	4.2						3.8
19	240	3.4						3.5
20	240	3.2						3.6
21	240	3.0						3.4
22	260	2.8						3.4
23	280	2.6						3.8

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 40

Calcutta, India (22.6°N, 88.4°E)

November 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	(240)	(3.6)						3.2
01	(210)	(3.6)						
02	(225)	(3.2)						---
03	(210)	(3.6)						3.3
04	(210)	(3.2)						
05	(210)	(2.8)						
06	(220)	(3.3)						3.2
07	(210)	(6.8)				2.0		
08	210	7.8				2.5	3.1	
09	210	8.7				2.8		3.3
10	225	10.8				3.2		
11	(210)	(10.6)				3.4		
12	(210)	(11.0)				---		3.1
13	210	11.0				3.2		
14	210	(11.0)				3.1		
15	210	(11.0)				3.2		3.4
16	210	11.0				2.6	3.0	
17	200	10.8				---		
18	180	9.4						3.3
19	190	7.1						
20	210	5.5					3.0	
21	(210)	(5.2)						3.2
22	210	4.1						
23	220	3.6						

Time: 90.0°E.

Sweep: 0.5 Mc to 18.0 Mc in 10 minutes, semi-automatic operation.

Table 41

Bombay, India (19.0°N, 73.0°E)

November 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06								
07	300	5.7						3.1
08:30	330	7.1						3.0
09	330	7.6						2.9
10	360	8.4						2.8
11	390	10.1						2.7
12	390	10.7						2.6
13	390	11.2						2.6
14	400	11.6						2.6
15	420	11.8						2.6
16	400	11.7						2.6
17	390	11.2						2.6
18	360	10.8						2.8
19	360	10.1						2.9
20	330	8.4						2.9
21	300	7.0						3.0
22	300	5.3						3.1
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 42

Madras, India (13.0°N, 80.2°E)

November 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	300	5.7						3.0
07	360	7.1						2.8
08	360	7.9						2.7
09	390	8.2						2.6
10	420	8.1						2.6
11	420	8.4						2.5
12	420	8.8						2.5
13	420	9.0						2.5
14	420	9.2						2.5
15	420	9.7						2.6
16	390	9.8						2.6
17	390	10.3						2.6
18	390	9.0						2.6
19	360	8.1						2.7
20	345	7.6						2.8
21	330	6.8						2.9
22	300	6.2						3.1
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 43

November 1953

Tiruchy, India (10.8°N, 78.8°E)

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	330	4.8						3.0
07	360	6.7						2.8
08	390	7.6						2.6
09	420	7.5						2.5
10	420	7.5						2.5
11	420	7.7						2.5
12	450	8.4						2.4
13	450	8.4						2.5
14	450	8.6						2.4
15	450	8.8						2.4
16	450	> 8.6						2.4
17	420	8.5						2.5
18	420	8.5						2.5
19	420	8.0						2.6
20	420	7.6						2.6
21	360	7.4						2.7
22								
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 44*

November 1953

Singapore, British Malaya (1.3°N, 103.8°E)

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	235	4.6					2.0	3.1
01	250	3.8					2.2	3.0
02	260	3.6					1.8	3.1
03	255	2.9					2.1	3.1
04	255	2.5					2.4	3.1
05	260	2.3					2.8	3.2
06	250	3.9			140	1.3	2.8	3.2
07	255	6.3	235		120	2.3	3.2	3.1
08	300	7.2	220	4.2	115	2.8	3.8	2.8
09	355	7.6	215	4.4	115	3.1	4.3	2.4
10	385	6.1	210	4.4	110	3.3	4.3	2.3
11	395	8.4	205	4.5	110	3.4	5.9	2.2
12	410	8.2	205	4.5	110	3.4	5.6	2.1
13	395	8.4	200	4.4	110	3.3	5.8	2.1
14	375	8.6	200	4.4	110	3.2	5.4	2.3
15	350	9.0	205	4.3	115	3.0	5.9	2.3
16	335	9.2	230	4.2	115	2.6	5.5	2.4
17	260	9.2			125	2.1	5.8	2.5
18	265	9.0				(1.4)	4.6	2.6
19	290	8.3					3.8	2.7
20	295	7.5					3.7	2.7
21	275	7.6					3.1	2.9
22	235	8.3					2.9	3.3
23	210	5.9					2.8	3.3

Time: 105.0°E.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 45

November 1953

Barotonga I. (21.3°S, 159.8°W)

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	270	6.2					3.0	3.05
01	250	6.4					2.8	3.1
02	250	5.3					3.0	3.0
03	285	4.5					2.6	2.9
04	300	4.2					2.4	2.9
05	290	4.2					2.3	3.0
06	250	5.9			1.9		2.9	3.2
07	250	6.7	230	3.8	110	2.5	3.5	3.3
08	280	7.3	220	4.2	110	2.8	4.1	3.2
09	300	7.6	210	4.4	105	3.1	4.5	3.05
10	310	8.4	200	4.5	105	3.3	4.6	3.0
11	310	9.4	200	4.5	105	3.4	4.5	3.0
12	305	10.3	200	4.5	105	3.4	4.2	3.0
13	310	10.3	200	4.5	105	3.4	4.2	3.0
14	290	10.5	200	4.5	105	3.3	4.1	3.1
15	290	10.1	210	4.4	110	3.2	3.8	3.1
16	280	9.6	220	4.1	110	2.9	3.9	3.1
17	280	9.1	240	---	110	2.5	4.1	3.1
18	260	8.7			---		4.0	3.1
19	260	7.8			---		4.0	3.0
20	280	7.1					4.0	2.8
21	300	6.9					3.8	2.8
22	300	6.6					3.2	2.8
23	300	6.4					3.0	2.8

Time: 157.5°W.

Sweep: 2.0 Mc to 16.0 Mc, manual operation.

Table 46*

October 1953

Inverness, Scotland (57.4°N, 4.2°W)

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	325	(1.9)						2.8
01	340	1.9						2.8
02	325	(1.8)						2.8
03	325	1.9						2.8
04	310	2.0					2.3	(2.9)
05	290	1.8					2.5	(3.0)
06	(265)	(2.0)					2.4	(3.1)
07	255	2.8			(150)	(1.6)	1.8	3.2
08	245	4.0	235	(3.0)	125	1.8	2.3	3.4
09	265	4.5	225	3.4	115	2.1	2.4	3.4
10	290	4.9	215	3.6	110	2.3	2.5	3.3
11	285	5.2	210	3.7	110	2.4	2.6	3.3
12	275	5.6	210	3.8	110	2.5	2.7	3.3
13	270	5.8	215	3.7	110	2.5	2.6	3.4
14	260	5.7	220	3.6	115	2.4	2.3	3.3
15	260	5.6	230	3.5	120	2.2	1.8	3.3
16	260	5.3	245		125	2.0		3.3
17	250	5.1			(145)	(1.6)	1.9	3.3
18	245	5.1						3.2
19	260	4.3						3.2
20	255	3.6						3.3
21	265	3.0						3.1
22	305	2.2						2.9
23	325	1.9						2.9

Time: 0.0°.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 47*

October 1953

Slough, England (51.5°N, 0.6°W)

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	295	2.9					2.4	2.8
01	285	2.9					2.5	2.8
02	285	2.9					2.6	2.8
03	285	2.9					2.6	2.8
04	280	2.6					2.6	3.0
05	265	2.4					3.2	3.0
06	270	2.4					2.5	3.0
07	250	3.9	(230)	(2.6)	135	1.7	2.7	3.3
08	265	4.8	235	3.2	125	2.1	3.4	3.4
09	280	5.4	230	3.6	120	2.4	3.8	3.3
10	280	5.8	225	3.8	120	2.6	4.0	3.4
11	280	6.1	215	3.9	120	2.7	3.9	3.4
12	280	6.3	220	3.9	120	2.8	3.9	3.4
13	260	6.0	225	3.8	120	2.7	3.9	3.4
14	260	6.1	230	3.7	120	2.5	3.6	3.4
15	260	6.1	240	3.5	125	2.3	3.5	3.3
16	250	5.8	235	3.2	130	2.0	2.9	3.4
17	240	5.5			(130)	(1.6)	3.1	3.4
18	245	5.3					3.1	3.2
19	245	4.4					2.6	3.2
20	245	3.7					2.4	3.2
21	270	3.0					2.6	3.0
22	285	2.9					2.4	2.9
23	290	2.9					2.4	2.8

Time: 0.0°.

Sweep: 0.55 Mc to 16.5 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 48

October 1953

Delhi, India (28.6°N, 77.1°E)

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	285	3.1						3.2
01	280	3.0						3.2
02	(280)	2.9						(3.3)
03								
04	280	3.2						3.3
05	260	3.3						3.4
06	240	4.1						3.5
07	240	6.7						3.6
08	240	7.4						3.6
09	240	7.9						3.6
10	240	8.1						3.4
11	280	8.2						3.3
12	280	9.1						3.2
13	280	9.4						3.3
14	260	> 9.4						3.4
15	260	> 9.4						3.4
16	240	8.1						3.4
17	240	7.9						3.6
18	240	6.8						3.6
19	240	4.2						3.6
20	240	4.0						3.4
21	260	> 3.5						3.4
22	280	3.2						3.3
23	290	3.4						3.1

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 49

Calcutta, India (22.6°N, 88.4°E)

October 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	(225)	4.2						3.1
01	(210)	3.6						
02	(240)	3.8						
03	(240)	3.8						3.1
04	(240)	3.1						
05	(210)	2.8						
06	210	4.6						3.3
07	210	8.5				2.3	3.1	
08	210	8.8				2.6	3.8	
09	210	9.2				2.9	3.4	3.1
10	225	10.2				3.2		
11	240	11.0				3.5		
12	240	11.0				3.8		3.1
13	225	11.0				3.6		
14	210	11.2				3.5		
15	210	11.0				3.1		3.3
16	210	11.3				2.6	3.0	
17	180	11.0				---	2.8	
18	180	10.8						3.5
19	180	8.8						
20	195	6.4						
21	(195)	5.0						3.35
22	(210)	4.9						
23	(230)	4.2						

Time: 90.0°E.

Sweep: 0.5 Mc to 19.0 Mc in 10 minutes, semi-automatic operation.

Table 50

Bombay, India (19.0°N, 73.0°E)

October 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06								
07	300	6.4						3.1
08	330	7.5						3.1
09	360	8.1						2.8
10	360	9.2						2.8
11	390	10.2						2.7
12	390	10.8						2.6
13	420	11.4						2.6
14	420	11.8						2.5
15	420	12.1						2.5
16	420	12.4						2.5
17	390	11.7						2.6
18	390	11.1						2.7
19	360	10.1						2.8
20	330	8.2						3.0
21	300	6.7						3.1
22	300	5.3						3.1
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 51*

Khartoum, Sudan (15.6°N, 32.6°E)

October 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	(275)	(6.5)					3.1	
01	(250)	(7.6)					3.1	
02	(225)	7.1					3.1	(3.3)
03	(215)	4.0					3.1	(3.3)
04	(230)	2.6					3.1	(3.3)
05	(265)	2.3					3.9	
06	(250)	4.7					3.4	(3.2)
07	(240)	6.9				(2.4)	5.2	(3.1)
08	(290)	8.3	(235)	(4.1)	(125)	(2.8)	4.1	(2.9)
09	(290)	10.0	(250)	(4.2)		(3.1)	4.6	(2.9)
10	(300)	10.4	(215)	(4.4)			4.8	(2.7)
11	(315)	10.5	(205)	(4.4)			4.6	(2.6)
12	(310)	10.8	(210)	(4.4)			6.9	(2.7)
13	(290)	11.5	(210)	(4.3)			6.9	(2.9)
14	(285)	11.8	(210)	(4.2)			6.9	(3.0)
15	(275)	12.0	(215)	(3.9)		(2.9)	6.8	(3.0)
16	(260)	11.7	(235)	(3.7)			6.4	(3.1)
17	(245)	11.6					5.7	(3.1)
18	(240)	10.8					4.4	(3.0)
19	(240)	9.6					3.1	(3.0)
20	(250)	8.7					4.0	(3.0)
21	(265)	7.9					6.2	(2.8)
22	(280)	6.9					6.2	(2.8)
23	(280)	(7.2)					3.1	

Time: 30.0°E.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 52

Madras, India (13.0°N, 80.2°E)

October 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	300	6.2						3.0
07	360	7.5						2.9
08	390	8.2						2.7
09	420	8.9						2.6
10	420	8.5						2.5
11	420	8.5						2.5
12	420	8.6						2.5
13	420	8.8						2.5
14	420	10.0						2.6
15	420	10.2						2.6
16	390	10.6						2.6
17	390	10.5						2.6
18	390	10.4						2.6
19	360	9.1						2.8
20	360	8.5						2.8
21	345	>7.6						2.9
22	(330)	---						(3.0)
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 53

Tiruchy, India (10.8°N, 78.8°E)

October 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	360	5.6						2.8
07	390	7.5						2.6
08	420	8.2						2.5
09	450	8.2						2.4
10	450	7.8						2.4
11	480	7.6						2.4
12	480	8.0						2.3
13	480	8.5						2.3
14	490	9.2						2.3
15	480	>9.8						2.4
16	450	10.5						2.4
17	420	>10.5						2.4
18	420	>9.5						2.5
19	420	>9.0						2.5
20	390	>8.7						2.6
21	360	8.3						2.7
22								
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 54*

Singapore, British Malaya (1.3°N, 103.8°E)

October 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	215	5.3					2.6	3.3
01	250	4.6					1.7	2.9
02	255	4.4					1.9	3.1
03	260	3.5					3.0	3.1
04	255	2.8					2.6	3.1
05	260	2.6					3.0	3.2
06	250	4.4			(140)	(1.3)	3.1	3.3
07	260	7.0	235		125	2.3	3.6	3.3
08	295	7.9	225		120	2.8	4.9	3.0
09	325	8.1	220	4.4	115	3.1	4.9	2.6
10	355	8.7	210	4.5	110	3.4	4.9	2.4
11	355	9.7	205	4.6	110	3.5	5.6	2.2
12	355	9.4	200	4.6	110	3.5	6.0	2.0
13	365	9.4	200	4.6	110	3.4	5.2	2.1
14	340	9.6	200	4.5	110	3.3	5.2	2.4
15	310	9.8	215		115	3.0	5.5	2.6
16	290	9.8	230		115	2.7	4.4	2.6
17	265	9.7	240		125	2.1	4.6	2.6
18	265	9.7					3.5	2.6
19	285	9.4					3.7	2.7
20	285	8.9					3.7	2.8
21	255	9.0					3.2	3.0
22	225	9.0					2.1	3.3
23	215	7.5					2.8	3.4

Time: 105.0°E.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 55

Nairobi, Kenya (1.3°S, 36.8°E) October 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	200	6.1						3.7
01	210	4.3						2.9
02	260	4.3						2.9
03	260	4.4					2.1	2.9
04	260	4.4					2.0	3.1
05	230	4.0					2.6	3.2
06	230	3.7					2.6	3.3
07	240	6.2	230	---	120	2.2	3.0	3.5
08	260	7.6	220	4.0	110	2.7	3.7	3.4
09	280	7.8	220	4.4	110	3.0	4.0	3.3
10	300	8.7	210	4.5	110	3.3	3.7	3.0
11	310	9.4	---	4.7	100	---	4.0	3.0
12	320	10.2	---	4.7	110	---		2.9
13	330	10.0	---	4.7	100	---		2.9
14	350	10.9	---	4.6	110	---	(4.7)	2.8
15	340	10.5	---	4.5	110	---	4.6	2.9
16	320	10.8	200	4.4	110	2.9	4.2	2.9
17	290	10.2	230	---	110	---	3.8	2.9
18	280	9.9	250	---	---	---	3.1	2.9
19	280	V 10.0					2.8	2.9
20	270	V 10.0					2.6	2.9
21	250	10.2						3.1
22	220	10.7						3.4
23	210	10.5						3.6

Time: 45.0°E.

Sweep: 1.0 Mc to 15.0 Mc in 7 seconds.

Table 56

Sao Paulo, Brazil (23.5°S, 46.5°W) October 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	250	7.8						3.2
01	240	7.2						3.3
02	220	8.4						3.4
03	220	5.8						3.3
04	220	4.8						3.3
05	220	4.3						3.5
06	210	5.5						3.8
07	220	6.6						3.6
08	260	7.0					3.3	3.4
09	280	7.5					3.2	3.0
10	310	8.4						2.9
11	355	9.4					4.0	2.8
12	320	10.0					4.0	3.0
13	325	10.4						3.0
14	300	11.8					4.0	3.1
15	280	12.2					3.3	3.2
16	260	12.4					3.4	3.4
17	240	12.5					3.2	3.4
18	220	12.4						3.5
19	210	11.6						(3.7)
20	200	10.0						3.4
21	220	3.9						3.3
22	230	9.0						3.5
23	240	8.7						3.1

Time: Local.

Sweep: 2.5 Mc to 20.0 Mc in 6 minutes.

Table 57*

Falkland Is. (51.7°S, 57.8°W) October 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	285	4.8						2.8
01	290	4.6						2.8
02	290	4.6						2.8
03	270	4.4						2.9
04	265	4.2						2.9
05	235	4.6						3.1
06	230	5.2	(240)					3.5
07	235	5.2	225		140	(1.4)		3.5
08	(290)	5.6	225	(4.0)	110	2.7	3.2	3.3
09	285	6.0	220	4.2	105	2.9	3.8	3.3
10	280	6.3	220	4.4	105	(3.0)	3.6	3.2
11	290	7.1	(205)	4.4	105	3.1	3.8	3.1
12	290	7.4	(215)	4.4	105	3.1	3.5	3.2
13	270	7.4	220	4.4	105	(3.1)	3.7	3.3
14	270	6.9	220	4.2	105	3.0	3.5	3.3
15	(260)	6.2	215	4.1	105	2.8	3.0	3.4
16	(255)	6.0	225	(3.8)	110	2.5	3.0	3.4
17	245	5.9	230		115	2.1	2.8	3.4
18	240	5.7			145	1.7	1.9	3.3
19	250	5.8					1.7	3.1
20	265	5.7						2.9
21	260	5.6						2.9
22	270	5.4						2.8
23	280	5.1						2.8

Time: 60.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average values except foF2 and fEs, which are median values.

Table 58*

Port Lockroy (64.8°S, 63.5°W) October 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	270	5.0						2.8
01	270	4.5						2.9
02	265	4.1						2.8
03	275	3.9						2.9
04	265	3.8				(135)	(1.4)	2.9
05	250	4.2			110	1.6	1.3	3.0
06	240	4.4			110	1.8		3.2
07	235	4.8			105	2.1		3.3
08	235	4.9			100	2.4		3.3
09	225	5.2			100	2.6		3.4
10	235	5.2	(210)		100	2.7		3.3
11	235	5.7	210		100	2.6		3.2
12	250	6.0	210	(3.8)	100	2.8		3.4
13	255	5.7	205	(4.0)	100	2.8		3.2
14	255	5.9	210	(4.0)	100	2.7		3.4
15	230	5.5	(210)		100	2.6		3.3
16	230	5.4			100	2.4		3.3
17	235	5.2			105	2.2		3.3
18	240	5.2			115	1.8		3.2
19	250	5.5			130	1.6		3.0
20	265	6.5						2.9
21	260	6.3						2.9
22	265	6.0						2.8
23	270	5.4						2.9

Time: 60.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes, automatic operation.

*Average values except foF2 and fEs, which are median values.

Table 59

Delhi, India (28.6°N, 77.1°E) September 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	(300)	3.8						(3.2)
01	(300)	3.2						(3.0)
02	(300)	(3.3)						(3.1)
03	---	---						---
04	280	3.2						3.2
05	260	3.8						3.4
06	240	4.7						3.6
07	240	6.0						3.5
08	240	7.1						3.5
09	240	7.0						3.4
10	280	7.0						3.2
11	320	8.0						3.1
12	280	9.1						3.2
13	280	9.4						3.2
14	280	9.0						3.2
15	280	8.9						3.3
16	260	8.1						3.4
17	260	7.5						3.4
18	240	7.0						3.4
19	240	6.0						3.6
20	260	4.6						3.4
21	280	3.8						3.2
22	310	3.6						3.1
23	300	3.2						3.1

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 60

Calcutta, India (22.6°N, 88.4°E) September 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	240	4.1						2.95
01	(240)	(4.4)						
02	(240)	(3.9)						
03	(240)	(3.6)						2.9
04	(240)	(3.0)						
05	(240)	(2.6)						
06	(210)	(4.5)						3.2
07	225	6.6						
08	225	8.2						
09	240	8.5						
10	240	10.2						
11	250	11.0						
12	255	10.3						
13	240	11.0						2.85
14	240	11.0						
15	255	11.0						
16	240	11.0						
17	210	11.0						
18	240	11.0						
19	210	10.1						
20	210	7.2						
21	210	5.4						
22	(240)	4.3						
23	(240)	4.3						3.2

Time: 90.0°E.

Sweep: 0.5 Mc to 19.0 Mc in 10 minutes, semi-automatic operation.

Table 61

Bombay, India (19.0°N, 73.0°E) September 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	(270)	(4.6)						(3.4)
07	270	6.4						3.2
08	300	7.4						3.0
09	330	7.9						2.9
10	360	8.7						2.8
11	390	9.8						2.6
12	420	10.6						2.6
13	420	11.0						2.6
14	420	11.6						2.5
15	420	11.7						2.5
16	420	11.9						2.6
17	390	11.9						2.6
18	390	11.4						2.8
19	330	9.8						3.0
20	330	7.6						3.0
21	300	6.1						3.0
22	300	4.6						3.1
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 62

Tiruchy, India (10.8°N, 78.8°E) September 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	360	5.1						2.9
07	420	7.2						2.6
08	450	7.8						2.4
09	480	7.4						2.4
10	480	7.2						2.3
11	480	7.5						2.3
12	480	7.4						2.3
13	480	7.8						2.3
14	480	8.4						2.3
15	480	9.6						2.4
16	450	9.9						2.4
17	450	10.2						2.4
18	450	9.8						2.4
19	435	8.5						2.5
20	420	>7.8						2.6
21	360	7.3						2.6
22								
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 63

Falkland Is. (51.7°S, 57.8°W) September 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	320	3.0						2.7
01	305	2.9						2.7
02	295	3.0						2.8
03	285	2.9						2.9
04	260	2.8						3.1
05	250	2.8						3.1
06	235	3.8			175	1.6	1.2	3.4
07	225	4.8			155	2.1	2.2	3.6
08	245	5.2	220		115	(2.4)	3.0	3.5
09	245	5.5	215		115	(2.5)	3.8	3.5
10	260	5.8	220	(4.0)	110	(2.8)	4.6	3.3
11	270	6.6	220	4.2	110	(2.9)	3.8	3.4
12	275	6.4	215	4.2	110	2.9	3.3	3.4
13	255	6.5	225	4.1	105	2.9	3.1	3.4
14	250	5.9	220	4.0	110	2.7	2.6	3.5
15	245	5.7	215	3.6	110	2.6	2.7	3.6
16	245	5.7	225		120	2.2	2.5	3.5
17	230	5.5			150	1.7	2.9	3.5
18	230	4.8						3.3
19	240	4.2						3.1
20	250	3.6						3.1
21	265	3.3						3.0
22	295	3.3						2.8
23	310	3.3						2.7

Time: 60.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes.

*Average value except foF2 and fEs, which are median values.

Table 64

Madras, India (13.0°N, 80.2°E) September 1953

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	330	5.9						3.0
07	360	7.5						2.8
08	375	8.1						2.6
09	420	8.2						2.6
10	420	8.1						2.6
11	420	8.0						2.5
12	420	8.5						2.5
13	420	8.8						2.4
14	420	9.2						2.4
15	420	9.3						2.5
16	420	10.4						2.6
17	420	10.8						2.6
18	375	10.4						2.6
19	360	>8.9						2.7
20	360	7.8						2.8
21	360	6.9						2.8
22								
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 65

Sao Paulo, Brazil (23.5°S, 46.5°W) September 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	230	7.0						3.4
01	220	5.6						3.5
02	225	5.8						3.7
03	220	4.0						3.4
04	240	3.5						(3.2)
05	260	3.2						(3.4)
06	232	3.8						3.5
07	220	5.7						3.6
08	240	6.4						3.4
09	280	7.8						3.3
10	300	8.6						3.2
11	290	9.7						3.0
12	300	9.8						3.1
13	300	10.0						3.2
14	280	11.0						3.2
15	260	10.9						3.5
16	220	10.3						3.6
17	220	9.0						3.5
18	205	9.2						3.6
19	200	(7.0)						(3.7)
20	200	6.6						(3.5)
21	220	6.6						3.3
22	235	6.8						3.2
23	240	5.2						3.4

Time: Local.

Sweep: 2.5 Mc to 20.0 Mc in 6 minutes.

Table 66

Fort Lockroy (64.8°S, 63.5°W) September 1953

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	2.7						2.7
01	290	2.7						2.7
02	290	2.6						2.8
03	290	2.5						2.8
04	270	2.5						2.9
05	260	2.4						3.0
06	240	3.0				(125)	(1.5)	3.1
07	235	4.0				125	1.8	3.4
08	225	4.5				(120)	(1.9)	3.5
09	230	5.0				110	2.3	3.4
10	230	5.2				105	2.4	3.4
11	220	5.2				(105)	(2.5)	3.3
12	220	5.5	(200)	(3.0)		(105)	(2.6)	3.4
13	225	5.6	(200)			(105)	(2.6)	3.3
14	230	5.5	(200)			105	2.4	3.4
15	230	5.5				110	2.2	3.4
16	230	5.1				115	2.0	3.4
17	235	5.0				(115)	(1.8)	3.3
18	235	4.8						3.3
19	240	4.6						3.1
20	250	4.4						3.0
21	265	3.7						2.8
22	280	3.1						2.7
23	300	3.0						2.8

Time: 60.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes, automatic operation.

*Average values except foF2 and fEs, which are median values.

Table 67

August 1953

Delhi, India (28.6°N, 77.1°E)

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								(3.1)
05	300	> 4.0						3.0
06	280	4.5						3.2
07	280	5.5						3.3
08	280	6.2						3.2
09	300	6.3						3.1
10	320	6.6						3.0
11	300	7.2						3.0
12	320	7.6						3.0
13	320	7.9						3.0
14	320	8.3						3.0
15	300	8.9						3.0
16	280	> 9.0						3.2
17	280	8.2						3.2
18	280	8.5						3.3
19	(280)	> 7.0						(3.4)
20	---	---						(3.2)
21	---	---						
22	---	---						
23	---	---						

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 68

August 1953

Bombay, India (19.0°N, 73.0°E)

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	270	4.8						3.4
07	300	5.9						3.2
08	330	7.2						3.0
09	330	7.4						2.9
10	380	8.2						2.7
11	420	8.9						2.6
12	420	9.4						2.5
13	420	10.1						2.5
14	420	10.4						2.5
15	420	10.9						2.6
16	360	11.3						2.7
17	330	10.8						2.8
18	330	10.3						2.9
19	320	9.3						3.0
20	300	7.8						3.1
21	300	6.2						3.0
22	300	5.1						3.2
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 69

August 1953

Madras, India (13.0°N, 80.2°E)

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	330	5.7						2.9
07	390	6.5						2.7
08	390	7.0						2.6
09	420	7.0						2.5
10	420	6.9						2.5
11	440	6.9						2.4
12	450	7.1						2.4
13	450	7.4						2.4
14	450	8.1						2.4
15	450	8.4						2.4
16	440	8.9						2.5
17	420	8.9						2.5
18	420	> 9.0						2.5
19	390	> 8.4						2.6
20	390	7.2						2.6
21	360	6.8						2.8
22								
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 70

August 1953

Tiruchy, India (10.8°N, 78.8°E)

Time	*	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00								
01								
02								
03								
04								
05								
06	360	4.6						2.9
07	390	6.4						2.6
08	420	6.8						2.4
09	460	6.9						2.4
10	480	6.6						2.3
11	500	6.7						2.3
12	510	6.8						2.2
13	510	7.0						2.2
14	510	7.3						2.2
15	480	> 7.8						2.2
16	480	8.5						2.3
17	480	8.6						2.4
18	480	8.4						2.4
19	450	7.9						2.5
20	420	7.0						2.6
21	390	6.0						2.6
22								
23								

Time: 75.0°E.

Sweep: 1.5 Mc to 18.0 Mc in 5 minutes, manual operation.

*Height at 0.83 foF2.

**Average values; other columns, median values.

Table 71

August 1953

Sao Paulo, Brazil (23.5°S, 46.5°W)

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	250	3.5						(3.3)
01	250	3.7						3.3
02	220	3.4						(3.4)
03	220	3.5						(3.7)
04	---	2.0						---
05	---	E						---
06	---	2.1						---
07	210	(4.4)						---
08	245	5.7						(3.5)
09	(300)	6.0						(3.4)
10	300	7.3						3.2
11	300	7.8						3.3
12	280	8.6						3.2
13	295	8.8						3.2
14	280	8.9						3.3
15	265	9.7						3.4
16	230	8.4						3.6
17	220	6.6						(3.4)
18	210	6.7						(3.6)
19	200	(3.6)						(3.5)
20	220	2.9						---
21	230	2.8						---
22	250	2.6						---
23	250	2.9						(3.3)

Time: Local.

Sweep: 2.5 Mc to 20.0 Mc in 6 minutes.

Table 72*

August 1953

Port Lockroy (64.8°S, 63.5°W)

Time	h'F2	foF2	h'F1	foF1	h'E	foE	fEs	(M3000)F2
00	300	2.2						2.9
01	290	2.2						2.9
02	290	2.2						2.9
03	280	2.3						2.9
04	260	2.2						3.1
05	240	2.0						3.2
06	235	1.8						3.2
07	235	1.8					1.3	3.2
08	240	3.0					1.4	3.3
09	220	3.9			(120)	(1.7)	2.0	3.6
10	220	4.2						3.5
11	220	4.8						3.6
12	220	4.5						3.6
13	225	4.6						3.6
14	225	4.4						3.6
15	225	4.5						3.5
16	225	4.2						3.5
17	220	3.8					1.4	3.4
18	230	3.1						3.2
19	250	2.4						3.2
20	265	2.0						3.0
21	285	2.0						2.9
22	300	2.1						(2.8)
23	305	2.1						2.8

Time: 60.0°W.

Sweep: 0.67 Mc to 25.0 Mc in 5 minutes, automatic operation.

*Average values except foF2 and fEs, which are median values.

TABLE 74

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

IONOSPHERIC DATA

Form expires June 1946

National Bureau of Standards
(Institution)Scaled by: E.J.W. J.W.P. J.J.S.Calculated by: E.J.W. J.W.P. J.J.S.

f^oF₂ _____ Mc _____ May _____ 1954
(Characteristic) (Unit) (Month)

Observed at Washington, D.C.
Lot 38.7°N, Long 77.1°W

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	(2.7)F	(2.5)F	(2.0)F	(2.2)F	2.2	2.4	3.3	3.7	(4.1)S	5.0	4.9	5.0	5.1	5.0	5.0	5.0	4.9	5.0	4.9	5.0	4.9	4.0	3.8	(3.3)F
2	3.1	2.8	2.6	2.4	1.9F	2.4	3.9	3.8	4.3	4.7	4.9	5.2M	5.4	5.1	5.3	5.2	4.8	5.3	5.8	5.3	5.4	4.5	3.9	3.3
3	2.9	2.6	2.4	2.4	2.4	2.6	3.7	4.4	4.7	4.9	5.4M	5.3	5.4	5.2	5.3	5.6	5.7	5.4	5.8	6.2	5.6	4.4	3.5	2.9F
4	2.4F	2.5F	2.4F	2.3F	2.4F	2.9F	4.2M	4.9	5.1	5.7M	5.4	5.8	5.8M	5.4	5.6	6.0	5.6	6.1	6.3	6.9	6.6	(5.4)F	2.9F	3.2
5	1.9	1.9F	(1.8)F	(1.7)F	1.8	2.2	3.3	3.4G	3.6G	4.2	4.4F	5.0	4.5	4.4	4.8	4.9	A	A	5.2F	5.5	4.9	[3.7]A	[2.6]A	
6	2.1F	(2.3)F	2.2F	2.0	2.1F	3.5F	3.7F	4.1	4.2	4.5	(4.7)F	4.9	4.7	4.6	5.0	5.4	5.0	4.9	4.8	5.0	4.8	4.1	2.8	2.7
7	2.4F	2.3	2.2	2.1	2.1F	3.0	4.2	4.5	4.9	(4.8)F	4.9	4.9	5.0	4.8	5.0	5.1	5.4	5.3	5.4	5.6	5.5	4.5	(3.7)F	3.6
8	3.2	3.0F	2.7F	2.4	2.3	2.9	3.8	4.4M	4.4M	4.6M	4.6	5.6	5.0	5.0	5.0	4.9	5.4	6.2	6.8	7.2	6.4	5.8	4.8	3.8
9	3.4F	2.6F	2.4	(2.1)F	(1.9)F	2.4	3.4M	3.8M	4.1	3.9G	3.9G	4.0F	4.4M	4.4M	4.4M	4.6M	4.5M	4.8M	4.9M	5.2M	4.8M	3.3M	2.5M	2.3M
10	K(2.1)F	(2.0)F	(1.9)F	(2.0)F	2.0F	2.9	4.1	4.9	4.9	5.0	5.1	5.3	5.4	5.0	5.2F	5.1	5.2	5.6	6.6	7.0	(6.1)F	4.6	3.8	3.5
11	3.0	2.7	2.5	2.3	1.9	2.4	3.2	3.7	4.1	4.3	4.5	4.7	4.9	4.7	5.0	5.0	5.5	5.2	6.1	5.9	5.2	5.0	3.4	2.5
12	2.9F	2.4	[2.2]A	1.9	[2.0]A	(2.6)F	3.6	4.2	4.4	5.4M	5.0M	4.6	4.9	5.1	5.0	5.1	5.2	[5.3]A	5.4	5.6	5.7	4.8	3.8	2.9F
13	2.5F	[2.9]A	2.8	2.3F	1.9F	A	A	A	4.3	[4.3]A	4.9M	4.8	4.5	[4.6]A	4.7	[4.9]A	5.1	5.6	[6.0]A	5.9	5.6	4.3	[3.4]A	[3.0]A
14	2.7F	2.5F	2.3F	2.3F	1.9	(2.0)F	A	A	3.5G	A	K	4.0F	4.3M	4.1F	4.2M	4.4M	4.5M	[4.6]A	[4.7]A	[4.7]A	5.0M	4.2F	3.7F	3.4M
15	3.1M	2.5F	2.3F	2.3F	(2.0)F	2.8	3.6M	4.4M	4.4	4.7	4.9M	4.9	4.9	4.9	4.8	4.9	4.5	5.3	5.8	6.5	6.1	4.8	4.2	3.6
16	2.9	2.1	1.9	1.8	1.8F	2.5	3.3G	3.9	4.3	(4.3)F	5.3	5.0	5.0	5.4M	5.3	4.9	4.9	5.2	5.6	6.2	6.0	[4.6]A	3.3	[2.8]A
17	2.4	2.2	[2.0]A	1.8	1.7	2.9	[4.3]A	5.3	5.0	5.0	5.2	5.2	5.3	5.0	5.2	5.3	5.2	5.4	5.4	6.0	(6.0)F	5.4	4.0F	3.2
18	3.0	2.9F	2.8	[2.4]A	2.3F	2.8	3.6	T	A	5.0	4.7	5.1M	4.5	4.6	4.8	5.1	5.0	5.2	5.5	A	A	5.6	(3.6)F	2.4F
19	2.2	2.1F	2.4	[2.3]A	2.2	2.6F	3.3	3.8	4.1	[4.3]A	(4.5)F	4.7	4.9	5.0	5.5M	5.4	5.0	5.4	5.6	6.5	(6.1)F	5.6	4.6F	4.0
20	3.1	2.8F	(2.5)F	(2.1)F	(2.0)F	2.5	3.3	3.9	4.2	(4.5)F	4.6	4.6	4.2M	4.8	4.7	4.9	5.4	6.3	6.6	(6.1)F	5.5	4.5	(3.5)F	(3.1)F
21	(3.1)F	2.8F	(2.0)F	5	S	2.5	3.3	3.6	4.3	(4.3)F	4.0F	4.6	4.9	5.1	4.6M	4.8	5.4	[5.9]A	6.4	7.6	5.6	3.3	2.8	2.3
22	2.3	(2.3)F	(2.1)F	1.9	1.9	2.9	3.4	3.9	4.2	5.0	4.3	4.6	4.8M	4.6M	5.1	4.9	5.1	5.8	(6.1)F	(6.4)F	4.8	3.9	3.2	
23	3.0	2.4	(2.0)F	(1.8)F	(1.9)F	2.6	3.4	4.1	5.0	4.5	[4.3]A	[4.4]A	4.6	[4.8]A	4.9M	4.9M	4.7	M	M	M	M	3.7	3.2	2.6
24	2.5	[2.4]A	(2.1)F	(1.9)F	(1.9)F	2.8	3.4M	3.6	(4.2)F	4.7M	4.7	[4.5]A	4.3	4.5	4.3	4.8	4.9	5.1	5.2	5.2	5.0	4.0	[3.4]A	(2.6)F
25	(2.5)F	2.5F	2.5	2.3F	2.3	3.2	4.1	4.3	4.9F	5.0M	5.6	[5.2]A	4.9	4.9	4.8	4.7	4.8	5.4	[5.9]A	6.0	5.8	4.8	3.6	3.1
26	2.9	A	A	A	A	A	3.5	[4.4]A	[5.0]A	[5.3]A	(5.1)F	4.9	4.8	4.7	(4.6)A	[5.0]A	[5.3]A	[5.5]A	5.8	[5.8]A	(5.1)F	(4.4)F	3.8	3.3
27	2.9F	2.6F	(2.2)F	(1.9)F	1.8F	2.6	3.9M	3.8	4.5	A	A	5.0	5.0	4.6	4.8	4.6F	4.8	[5.1]A	[5.8]A	6.6	6.0	4.6	3.1	2.6F
28	(2.3)F	2.3F	(2.2)F	A	A	3.3	4.3	4.5	5.0M	5.0F	5.4	[5.3]A	5.2	4.8	4.7	4.9	5.3	6.2	6.2	(6.8)M	6.0	4.5F	4.2	3.4F
29	3.2F	2.9F	2.4	2.0F	2.2F	(2.5)F	3.3	4.0	4.2	4.8	4.9	4.7	4.6	4.5	4.7	4.8	4.9	4.9	5.3	5.8	5.8	4.9	3.8	3.1
30	3.1	2.8	2.4	(2.2)F	2.0	3.0	3.9	4.4	4.9	4.9M	5.3	5.1	5.0	5.2	4.7	5.0	5.0	5.1	5.7	6.1F	6.2	5.2	3.7	3.0
31	2.7	2.6	2.5F	2.3F	(2.0)F	3.0M	3.5M	4.9	5.5	5.0	4.9	5.0	4.9	4.7	(4.4)F	4.5	4.9	5.3	[6.0]M	6.4	6.3	5.0	3.7	3.0
Median	2.4	2.5	2.3	2.2	2.0	2.7	3.6	4.1	4.4	4.8	4.9	4.9	4.9	4.8	4.8	4.9	5.0	5.3	5.8	6.0	5.7	4.6	3.7	3.0
Count	31	30	30	28	28	28	29	29	29	29	29	31	31	31	31	31	30	29	30	29	29	31	31	31

Sweep 1.0—Mc 1625.0—Mc in 0.25—min

Manual ☐ Automatic ☒

TABLE 75

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

National Bureau of Standards
(Institution)

Scaled by: E.W. J.W.P. J.J.S.

Observed at

Washington, D.C.

Lot 38.7°N, Long 77.1°W

f°F2 Mc M30 54

(Unit) (Month)

75°W Mean Time

Day	0030	0130	0230	0330	0430	0530	0630	0730	0830	0930	1030	1130	1230	1330	1430	1530	1630	1730	1830	1930	2030	2130	2230	2330
1	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°	(0.5)°
2	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3
3	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4	4.5	4.6	4.7	4.8	4.9	5.0	5.1	5.2	5.3	5.4
4	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°	(2.5)°
5	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4
6	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
7	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
8	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
9	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
10	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
11	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
12	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
13	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
14	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
15	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
16	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
17	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
18	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
19	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
20	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
21	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
22	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
23	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
24	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
25	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
26	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
27	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
28	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
29	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
30	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
31	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4
Median																								
Count																								

Sweep 10 — Mc 1025.0 — Mc 1025.5 min

Manual ☐ Automatic ☒

Form adopted June 1946

TABLE 76
Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

h'F1 km May, 1954
(Characteristic) (Unit) (North)

Observed at Washington, D. C.

IONOSPHERIC DATA

National Bureau of Standards
(Institution)

Scoted by: E.J.W., J.W.P., J.J.S.

Calculated by: E.J.W., J.W.P., J.J.S.

Lat. 38.7°N, Long 77.1°W

75°W Mean Time

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1							210	210	230	200 ^H	200	200 ^H	190	230	210	220	250	A	A					
2							210	210	210 ^H	230	200	210 ^H	220	230	220	220	200 ^H	190 ^H	230					
3							210	210	200 ^H	200 ^H	200 ^H	210 ^H	190 ^H	230	200 ^H	240	240 ^H	220	A					
4							240	230	240 ^H	230 ^H	210	200 ^H	190	200	190 ^H	240 ^H	240 ^H	260	250					
5							240	220	220 ^H	210	200	190	180	210 ^H	A	A	A	A	A					
6							210 ^H	200 ^H	200	200 ^H	250 ^H	250 ^H	210 ^H	210	190	A	A	A	200					
7							A	250 ^H	A	A	200 ^H	210 ^H	220	220 ^H	220	210 ^H	220 ^H	230	230					
8							220	200 ^H	200 ^H	200 ^H	200	200 ^H	220	210	210	200 ^H	240 ^H	240	240					
9							250	230 ^H	200 ^H	200 ^H	210 ^H	200 ^H	180 ^H	220 ^H	220 ^H	340 ^H	240 ^H	A ^K	A ^K					
10							220	210 ^H	210	200 ^H	200 ^H	200 ^H	200	190 ^H	210	210	230	A	A					
11							220	220	200 ^H	210	210 ^H	200	220 ^H	240	240 ^H	A	A	A	A					
12							A	230	220 ^H	220	230 ^H	210 ^H	A	A	A	A	A	A	A					
13							A	A	A	A	210	190 ^H	190 ^H	200 ^H	220 ^H	A	A	A	A					
14							A	220 ^H	A	A	200 ^H	200 ^H	190 ^H	200 ^H	210 ^H	220 ^H	A ^K	A ^K	A ^K					
15							210 ^H	180 ^H	210 ^H	210	190 ^H	200	190	200	210	220	220	230 ^H	A					
16							230	220 ^H	210 ^H	200	200 ^H	200 ^H	210 ^H	200 ^H	190 ^H	210	210 ^H	210 ^H	A					
17							A	200	200	200 ^H	A	210 ^H	190	190	220	220 ^H	220 ^H	210 ^H	A					
18							A	T	A	200	200	200	210	190	200 ^H	190 ^H	220	210	210 ^H					
19							220	220 ^H	A	A	A	A	190	210	220 ^H	210	200 ^H	180 ^H	240					
20							220	210	210	200	220	180 ^H	200	200 ^H	210	200 ^H	240	230	A					
21							220	210	220 ^H	190 ^H	180 ^H	200	200 ^H	190	200	220	210	250	A					
22							210	200	210	200 ^H	190	190 ^H	210	190	220	A	A	230 ^H	A					
23							220	200 ^H	200 ^H	210 ^H	A	A	170 ^H	A	A	A	A	M	M					
24							220	210	210	200 ^H	170 ^H	190 ^H	190 ^H	200 ^H	220	220	230 ^H	240 ^H	210					
25							A	210 ^H	210 ^H	180 ^H	200	200 ^H	200	180 ^H	200 ^H	220	A	A	A					
26							220	A	A	A	200 ^H	200 ^H	200	A	A	A	A	A	A					
27							210	210	A	A	A	A	200 ^H	200 ^H	200	210	A	A	A					
28							220	220	A	A	A	A	170 ^H	170 ^H	180 ^H	220	220	A	A					
29							280 ^H	230 ^H	220 ^H	210	210	210 ^H	210 ^H	220 ^H	200 ^H	230	220	240	A					
30							230	210	210 ^H	210 ^H	200 ^H	200 ^H	210 ^H	200 ^H	200 ^H	230 ^H	230	A	A					
31							240 ^H	230	240 ^H	180 ^H	170 ^H	200 ^H	210 ^H	190 ^H	210 ^H	220	230	A	M					
Median							220	210	210	200	200	200	200	200	210	220	220	230	240					
Count							24	27	22	23	25	27	30	28	27	23	20	14	8					

Sweep 1.0 Mc to 25.0 Mc in 0.25 min
Manual ☐ Automatic ☒

TABLE 77

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

National Bureau of Standards

(Institution)

Scaled by: E.J.W. J.W.P. J.J.S.

Calculated by: E.J.W. J.W.P. J.J.S.

IONOSPHERIC DATA

Lat. 38.7°N, Long. 77.1°W

Observed at Washington, D.C.

f°F1 _____ Mc _____ May _____ 1954

(Unit)

(Month)

75°W Mean Time

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	L						L	3.4	3.7	4.0	4.1	4.2	4.1	4.2	4.2	4.0	3.8	A	A					
2	L						L	3.4	3.7	4.0	4.1	4.2	4.1	4.2	4.2	4.0	3.8	3.6	3.6					
3	L						L	3.5	3.7	4.1	4.1	4.1	4.2	4.2	4.1	4.1	3.8	3.5	L					
4	L						L	3.5	3.7	4.2	4.2	4.1	4.2	4.2	4.1	4.1	3.8	3.6	3.1					
5	L						L	3.4	3.6	4.1	4.1	4.1	4.1	4.1	4.0	4.0	4.1	A	A					
6	L						L	3.5	3.7	4.0	4.1	4.2	4.2	4.1	4.0	4.0	3.8	3.8	3.0					
7	A						A	3.5	3.8	4.0	4.1	4.2	4.2	4.2	4.2	4.1	3.8	3.6	L					
8	L						L	3.6	3.7	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
9	L						L	3.4	3.7	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
10	L						L	3.6	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
11	L						L	3.4	3.6	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
12	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
13	A						A	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
14	A						A	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
15	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
16	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
17	A						A	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
18	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
19	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
20	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
21	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
22	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
23	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
24	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
25	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
26	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
27	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
28	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
29	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
30	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
31	L						L	3.5	3.8	4.3	4.3	4.3	4.3	4.3	4.3	4.0	3.8	3.6	3.1					
Median																								
Count																								

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

Manual ☐ Automatic ☒

Form expires June 1946

TABLE 78

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

h'E (Characteristic) Km (Unit) May, 1954
Observed at Washington, D.C.
Lat. 38.7°N, Long. 77.1°W

National Bureau of Standards
(Institution)
Scaled by: E.J.W., J.W.P., J.J.S.
Calculated by: E.J.W., J.W.P., J.J.S.

IONOSPHERIC DATA

75°W

Mean Time

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1							120	110	110	110	100	100	100	100	100	100	110	110	120					
2							S	110	110	110	100	100	(100) ^A	100	100	(100) ^A	110	110	110					
3							(110) ^S	110 ^H	110	110	110 ^H	100 ^H	100	100	100	100	100	110	S					
4							120 ^H	110	110 ^H	110	110	100	100	100	100	100 ^H	110 ^H	110 ^H	120					
5							(130) ^S	110 ^H	110 ^H	110	100	100	100	100	100	100 ^B	110 ^H	110	120					
6							A	110	110	110	(110) ^C	110	110	110	110	110	110	110	S					
7							S	110	110	110	100	100	100	100	100	100	110	110	(120) ^S					
8							110 ^H	110 ^H	110	110	100	100	100 ^H	110	110 ^H	110	110 ^H	110	(120) ^S					
9							S	110	110 ^H	110	100	100 ^K	100 ^K	100 ^K	100 ^K	100 ^K	110 ^K	110 ^K	120 ^K					
10							120	110	110	110	110	110	100	100	110	110	110	110	(110) ^S					
11							S	110	100	100	100	110	100 ^H	100	100	100	110	110	120					
12							(120) ^S	110	110 ^H	100 ^H	100 ^H	100 ^H	100 ^H	100 ^H	100	100	100	120	110					
13							(110) ^S	110 ^H	100	100	100	100	110	100	110 ^H	110	110	120	120					
14							(120) ^S	110 ^H	110	100 ^K	100 ^K	100 ^K	100 ^K	100 ^K	100 ^K	110 ^K	100 ^K	110 ^K	120 ^K					
15							A	A	100	100 ^H	100	100 ^H	100	100	100	110	110	110	120 ^H					
16							120	110	100	100	100	100	100	100	100 ^H	100	100	100	120					
17							120 ^H	110 ^H	100	100	100 ^H	100 ^H	A	A	A	100	100	100	120 ^S					
18							120 ^S	110 ^H	100	100	100	100	100	100	100	100	(110) ^B	110	(120) ^B					
19							110 ^S	110	100	100 ^H	100	100	100	100	100	100	110	110	110					
20							A	A	100 ^H	100 ^H	100	100	100 ^H	100	(110) ^B	(110) ^B	(110) ^B	110 ^H	(110) ^S					
21							(120) ^S	110 ^H	100	100	100	100 ^H	100 ^H	100 ^H	100	100	100	100	110					
22							S	110 ^H	100	100 ^H	(100) ^B	100	100	100	110	100	100	110	(100) ^S					
23							(120) ^S	100 ^H	100	100 ^H	100 ^H	100	100	(100) ^B	100 ^H	100	AA	M	M					
24							(110) ^S	110	100	100	100	100	100 ^H	100 ^H	100	100	110 ^B	110	110					
25							110	110	100	100	100	100	100	100	100	100	100	100	110					
26							110	100	100	100	100	100	100	100	100	100 ^H	110	110	110					
27							110	100	100	100	100	100	100	100	100	100	100	110	110					
28							110 ^H	100	100	100	100	100	100	100	100	100	100	110	110					
29							120	110	110	110	110	110	100	110	100	110	110	110	110					
30							120	110	110 ^H	100	100	100	100	100	100 ^H	100 ^H	110 ^H	110 ^H	110 ^H					
31							110 ^H	110	110	100	110	(100) ^B	100 ^H	110	110	110	110	110	M					
Median							120	110	100	100	100	100	100	100	100	100	110	110	110					
Count							23	29	31	31	31	31	30	30	30	31	30	30	27					

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

Manual ☐ Automatic ☒

TABLE 79

f°E _____ Mc _____ May _____ 1954
(Characteristic) (Unit) (Month)

Observed at **Washington, D.C.**

Lat. **38.7°N**, Long. **77.1°W**

National Bureau of Standards
Scaled by: **E.J.W. J.W.P. J.J.S.**
Calculated by: **E.J.W. J.W.P. J.J.S.**

75°W Mean Time

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1							1.7	2.3	2.7	2.8	2.8	(2.9) ^F	(3.0) ^A	3.2 ^H	A	A	B	2.5	S					
2							1.8 ^H	2.1 ^H	2.6	2.8	2.8	(3.2) ^A	3.2 ^H	3.2	(3.7) ^A	(3.0) ^S	(2.8) ^P	2.4	1.9					
3							1.9 ^H	2.4 ^H	2.8	2.9	3.1 ^H	3.1 ^H	3.2	3.1 ^H	(3.0) ^S	(2.9) ^A	(2.8) ^S	2.2	S					
4							1.8 ^H	2.3	2.6 ^H	2.9	3.0	3.0	3.1 ^A	3.1	3.0	2.1 ^H	2.6 ^H	2.4 ^H	1.9					
5							1.7	2.2 ^H	2.6 ^H	2.8	(3.0) ^A	A	3.0	2.4	(2.1) ^B	2.7 ^H	2.4	(1.8) ^P						
6							A	A	(2.6) ^A	2.7	(2.8) ^C	(3.5) ^H	(3.0) ^A	3.2	3.1	(2.9) ^P	2.7	2.4	A					
7							A	2.4	2.6	2.6	2.8	(2.9) ^H	A	A	3.0	A	2.8	2.4	2.0					
8							1.8 ^H	2.4 ^H	A	A	3.0	3.2	3.2 ^H	3.2	3.1 ^H	3.0	2.8 ^H	2.5	2.0					
9							A	2.1	2.6 ^H	(2.8) ^A	2.9	(3.5) ^H	(3.2) ^K	3.1 ^K	2.9 ^A	2.7 ^K	2.4 ^K	1.9 ^K						
10							1.8	(2.3) ^P	2.6	2.7	(2.8) ^A	A	A	A	3.1	3.0	2.7	2.4	(1.9) ^A					
11							S	2.2	2.5 ^S	2.7	3.0	3.0	3.2 ^H	3.2 ^H	3.0	2.9	2.7	2.4	1.8					
12							1.7	2.3	2.6 ^H	2.9 ^H	3.1 ^H	3.1 ^H	3.2 ^H	3.1	3.0	(2.9) ^A	2.1	2.5	1.9					
13							1.8	2.3 ^H	2.5	2.6	2.7	A ^K	A	A	3.3 ^H	3.1	2.8	2.4	1.9					
14							(1.8) ^A	2.3 ^H	2.6	A ^K	A ^K	A ^K	A ^K	A ^K	3.2 ^K	3.0 ^K	2.8 ^K	2.5	2.0 ^K					
15							A	A	2.6	2.8 ^H	3.0	3.1 ^H	3.2	(3.1) ^A	(2.9) ^A	(2.8) ^A	2.8	2.5	1.9 ^H					
16							1.8	2.4	2.7	2.9	2.8	3.1	3.0	3.2 ^H	(3.1) ^P	A	A	1.9						
17							1.9 ^H	2.4 ^H	2.6	2.8	2.9 ^H	3.1 ^H	A	A	A	3.0	2.8	2.3	(1.8) ^B					
18							1.8	(2.3) ^H	2.6	2.7	2.8	2.8	(3.3) ^A	(3.2) ^A	(3.1) ^A	3.0	(2.1) ^B	2.5	(2.0) ^B					
19							2.0	2.4	2.7	2.9 ^H	3.0	3.1	3.0	2.9	(2.7) ^P	2.5	(2.5) ^A	2.5	2.1 ^S					
20							A	A	2.7 ^H	2.8 ^H	(2.9) ^F	(2.9) ^B	3.0 ^H	(3.0) ^P	(3.1) ^P	3.0	(2.9) ^B	2.6 ^H	2.6					
21							(1.8) ^A	2.2 ^H	2.5	(2.8) ^A	A	A	(3.1) ^H	(3.0) ^B	3.0	2.8	2.8	2.5	A					
22							S	2.3 ^H	2.5	2.8 ^H	(3.0) ^A	3.0	(3.2) ^P	(3.1) ^A	(3.0) ^P	3.0	(2.7) ^A	(2.5) ^S	(2.1) ^H					
23							2.0 ^H	2.4 ^H	(2.6) ^A	2.7 ^H	2.9 ^H	3.0	(3.1) ^P	(3.1) ^B	3.1 ^H	3.0	M	M	M					
24							(1.8) ^H	A	A	(2.9) ^P	3.2	3.2	(3.2) ^P	(3.2) ^B	3.1 ^A	3.0 ^H	2.8	2.4	1.8					
25							1.9	2.3	2.6	A	A	A	A	A	A	3.0	2.9	2.5	2.0 ^H					
26							2.2 ^H	2.4	2.8	3.0	3.0	3.0	3.2	3.0	(3.0) ^A	3.0 ^H	2.8	2.5	2.0					
27							1.7	2.3	2.8	2.8	3.0	(3.0) ^A	(3.2) ^A	(3.2) ^A	3.2 ^H	2.9	(2.8) ^P	2.5	1.8					
28							1.8 ^H	2.3	2.5	2.8 ^H	A	A	A	3.1	3.2	3.1	2.9 ^H	2.5	1.8					
29							1.9 ^H	2.5	2.6	A	A	A	3.3	3.1	A	A	A	(2.6) ^P	1.9					
30							2.1	2.4	2.7 ^H	2.8	2.8	A	A	A	A	2.9 ^H	3.0 ^H	2.6 ^H	2.1 ^H					
31							1.9 ^H	2.4	2.7	2.8	3.0	3.2 ^H	3.2 ^H	(3.2) ^A	3.1	3.0	2.9	2.5 ^H	M					
Median																								
Count							1.8	2.3	2.6	2.8	3.0	3.0	3.2	3.2	3.1	3.0	2.8	2.5	1.9					

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

Manual ☐ Automatic ☒

TABLE 80

Control Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

IONOSPHERIC DATA

National Bureau of Standards

(Institution)

Scaled by: E.J.W. JWP, J.J.S.

Calculated by: E.J.W. JWP, J.J.S.

Es (Characteristic) Mc, Km (Unit) May 1954

Observed at Washington, D.C.

Lot 38.7°N, Long 77.1°W

75°W

Mean Time

Doy	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	E	E	E	E	2.7 120	2.5 120	3.7 110	2.6 130	4.1 120	3.1 110	4.1 110	4.4 110	3.4 100	G	3.3 110	4.4 110	4.2 110	5.0 120	6.8 120	3.6 120	5.8 120	3.8 110	3.5 110	E
2	3.2 110	4.6 110	3.4 110	2.7 110	2.9 110	E	2.9 110	4.7 110	4.0 110	4.3 110	4.3 110	4.1 110	3.9 100	G	3.4 110	3.8 110	4.8 110	3.6 110	G	3.3 100	4.3 100	4.1 100	E	E
3	E	3.4 100	E	3.3 100	3.9 100	E	G	G	G	4.0 120	4.4 110	3.5 110	G	G	G	3.8 100	G	3.1 110	3.9 120	3.3 120	4.8 100	4.1 120	3.3 120	
4	2.2 100	3.4 100	2.5 100	2.5 100	3.2 100	3.8 120	2.4 130	3.8 120	4.1 120	5.0 110	4.0 110	3.5 110	4.3 100	G	G	G	G	4.2 130	3.9 120	4.8 110	3.8 110	4.8 110	3.7 110	
5	E	E	3.6 130	3.1 130	2.2 120	4.3 120	3.1 130	3.9 120	3.7 120	1.0 120	3.2 120	4.4 110	4.2 110	G	5.6 110	4.5 110	8.4 120	7.6 120	8.0 120	9.8 110	5.4 110	7.8 110	10.5 110	6.8 100
6	4.0 100	2.9 110	E	3.0 110	E	E	3.8 110	8.8 110	4.7 110	4.3 110	5.4 110	5.4 110	4.5 110	4.4 110	G	3.5 120	4.5 120	4.3 120	3.3 120	E	E	E	E	2.4 120
7	2.9 120	2.4 120	4.1 110	3.3 110	2.6 120	4.2 110	3.8 110	5.8 110	4.5 110	4.8 110	4.2 110	5.0 110	4.5 110	4.9 100	3.6 110	3.1 100	G	3.9 120	3.3 120	2.0 120	4.7 110	4.2 110	2.8 110	2.7 110
8	E	2.5 110	E	2.9 110	3.3 100	6.6 110	1.9 120	4.5 100	3.4 110	3.8 110	G	G	G	G	G	G	G	G	G	E	E	E	E	E
9	E	E	E	E	E	E	1.8 110	3.2 120	3.3 120	3.2 110	3.2 120	G	G	G	G	G	3.3 130	4.2 120	4.5 120	3.0 110	E	E	E	E
10	E	E	E	E	4.7 120	2.8 110	3.2 110	G	3.7 120	2.9 120	7.0 110	4.7 110	3.1 100	3.2 110	G	G	4.0 130	4.3 120	4.7 120	3.7 110	E	E	E	E
11	E	E	E	E	E	E	2.1 120	3.6 120	7.2 110	2.9 110	G	3.1 110	3.4 120	G	3.7 130	4.9 120	5.5 120	5.0 120	4.9 110	5.8 110	3.5 110	3.1 110	3.0 110	2.8 110
12	E	E	3.8 120	3.3 120	4.0 110	3.5 110	3.7 120	4.0 120	5.8 110	5.2 110	3.3 110	4.7 110	4.8 110	12.5 110	10.6 110	5.8 110	6.3 120	7.0 120	5.6 110	9.0 100	4.3 100	2.3 110	4.3 100	3.8 100
13	4.5 100	4.7 100	3.7 100	3.2 110	7.6 110	6.6 100	7.6 110	7.0 110	7.2 100	6.6 100	6.6 100	4.2 110	3.7 110	6.3 110	G	6.0 130	7.0 120	8.4 110	7.8 110	5.0 110	5.0 110	6.8 100	6.4 100	8.4 100
14	5.4 100	2.9 100	2.9 100	2.8 100	3.5 110	4.0 110	4.2 110	7.2 100	10.5 110	11.2 100	12.7 100	4.6 100	3.2 110	4.4 110	10.5 110	6.8 110	5.5 120	6.9 110	5.8 110	7.4 110	5.8 110	5.1 110	4.5 100	5.0 100
15	2.7 100	E	E	E	2.9 120	3.2 110	3.0 110	3.4 100	G	3.5 110	G	G	G	3.5 110	3.3 110	3.8 110	3.2 130	3.9 120	4.8 110	5.8 110	4.2 110	3.5 110	3.7 110	2.3 110
16	E	E	E	E	3.3 120	2.9 120	G	G	4.6 110	4.8 100	4.5 110	3.3 110	3.2 110	G	G	4.3 100	4.3 100	4.9 110	5.1 110	5.4 110	4.1 110	7.6 110	6.8 100	7.0 100
17	4.1 100	3.9 100	4.0 100	3.1 100	3.9 100	3.1 120	7.0 110	4.5 110	5.0 110	6.6 110	5.5 110	5.2 110	3.5 100	5.4 100	4.2 100	6.3 110	4.1 110	3.9 110	3.7 120	4.0 110	3.8 110	3.6 110	E	4.3 110
18	3.0 110	2.8 110	5.0 100	5.0 100	3.4 120	2.9 130	4.0 120	T	7.0 110	4.2 110	4.1 110	3.4 110	4.0 110	3.9 110	4.3 100	G	G	G	3.8 120	8.6 110	7.2 110	5.2 110	5.0 110	7.0 110
19	6.8 100	4.8 100	4.0 100	3.9 100	1.7 100	E	3.1 110	G	4.4 120	4.8 110	5.0 110	4.6 110	3.6 110	3.4 110	3.9 110	3.9 110	3.0 110	G	3.8 130	3.2 120	4.5 110	4.8 110	4.2 110	3.3 110
20	3.0 100	4.0 100	5.2 110	E	4.1 100	3.0 100	2.0 110	3.3 110	3.8 120	4.4 110	G	G	G	G	G	G	G	3.0 140	4.7 120	4.8 110	3.0 110	3.0 110	2.4 110	E
21	E	E	E	2.5 110	S	E	3.3 120	3.4 110	3.1 110	3.1 120	3.0 120	3.1 100	G	G	G	G	G	7.0 120	4.2 120	5.5 110	4.4 110	4.3 110	E	2.8 110
22	E	E	E	3.0 140	3.1 120	3.1 110	3.0 120	2.7 120	7.6 100	4.4 110	4.3 110	3.8 110	G	4.1 110	3.4 120	5.6 120	8.0 120	3.9 130	5.5 120	5.0 110	3.9 110	E	E	E
23	E	E	E	E	2.3 110	E	E	3.5 110	4.3 110	5.7 110	5.3 110	5.4 110	G	5.0 140	7.0 130	6.6 120	5.6 120	M	M	M	M	E	2.9 110	3.7 110
24	E	3.8 120	7.4 120	7.3 110	5.0 110	2.9 120	3.3 120	4.1 110	5.6 110	5.0 110	4.3 120	10.5 110	7.8 110	5.0 130	G	G	4.8 110	7.0 110	3.7 110	3.7 110	4.4 110	3.1 100	5.0 100	4.4 100
25	2.5 110	E	E	2.9 120	4.3 110	3.0 110	5.2 110	5.2 100	3.6 110	4.3 100	7.4 100	8.4 100	4.4 100	3.6 100	4.7 100	5.2 120	4.5 120	5.5 110	6.8 110	5.0 110	6.0 110	5.4 100	2.5 110	3.0 110
26	2.9 110	7.4 100	4.4 100	6.7 100	5.8 100	4.3 100	3.4 120	5.8 110	7.4 110	6.2 110	5.3 110	5.6 110	5.4 110	7.8 110	7.0 100	9.0 110	7.6 110	1.3 110	8.0 110	9.8 110	3.8 110	4.2 110	3.9 110	3.0 110
27	4.0 100	7.2 100	7.6 110	E	E	2.6 110	3.9 110	3.8 110	5.0 110	6.7 100	8.0 100	6.6 100	7.4 100	5.9 100	G	4.5 120	5.2 120	7.4 110	1.5 110	7.0 100	3.9 100	2.5 100	E	3.3 100
28	5.8 100	3.9 100	5.4 100	4.2 100	3.7 100	4.0 100	3.0 120	5.3 100	7.7 100	7.4 100	6.0 100	12.5 100	4.0 100	3.4 110	G	G	4.0 110	4.4 110	5.4 110	8.8 110	7.0 110	3.8 110	3.1 110	2.7 110
29	2.4 110	5.2 110	4.2 110	E	3.0 110	E	2.9 120	4.8 120	4.0 110	4.4 110	5.0 110	4.5 110	4.9 110	G	4.0 100	3.9 110	3.7 110	G	3.8 120	3.2 110	3.2 110	2.5 110	4.3 110	5.7 110
30	3.5 110	4.3 110	E	4.1 100	3.0 100	E	7.6 100	3.6 120	4.2 110	4.2 110	5.0 110	3.4 100	4.2 100	4.5 100	5.4 100	G	G	4.2 120	4.0 120	3.8 110	3.0 110	E	E	E
31	3.5 100	E	E	3.2 120	3.6 120	6.0 110	3.5 120	3.7 120	6.8 110	5.4 110	G	G	G	4.4 100	6.4 120	G	G	4.4 120	M	3.7 120	5.8 110	E	E	E
Median	2.5	2.8	2.9	2.9	3.2	3.0	3.1	3.8	4.4	4.4	4.3	4.4	3.7	3.5	3.3	3.8	4.1	4.3	4.7	4.4	4.3	3.5	3.0	3.0
Count	31	31	31	31	30	31	31	30	31	31	30	31	31	31	31	31	31	31	29	30	30	31	31	31

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

Manual ☐ Automatic ☒

TABLE 81

(M1500)F2

(Characteristic)

May, 1954

(Unit)

Observed at Washington, D.C.

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

IONOSPHERIC DATA

National Bureau of Standards

(Institution)

Scaled by: E.J.W., J.W.P., J.J.S.

Calculated by: E.J.W., J.W.P., J.J.S.

		75°W												Mean Time											
		Long 77.10W												Lat. 38.7°N											
Day		00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	(1.9) ²	(1.9) ²	(1.9) ²	(2.0) ²	(2.3) ²	2.2	2.3	2.3	1.9	(1.9) ²	2.2	2.3	2.0	2.2	2.0	2.1	2.1	2.1	2.3	2.3	2.3	2.2	2.0	2.0	(2.1) ²
2	1.9	2.0	2.2	2.2	2.2 ⁵	2.2	2.3	2.3	2.4	2.0	1.9	2.1	1.8 ^M	2.3	2.1	2.2	2.1	2.1	2.2	2.3	2.2	2.2	2.1	2.2	2.1
3	2.1	2.0	2.1	2.2	2.3	2.4	2.4	2.4 ^M	2.4	2.4	2.1 ^M	2.2	2.2	2.4	2.1	2.1	2.2	2.2	2.2	2.3	2.3	2.3	(2.4) ²	2.3 ^F	2.0
4	2.2 ⁵	2.2 ⁵	2.2 ⁵	2.3 ^F	2.4 ^F	2.4 ^F	2.4 ^F	2.4 ^F	2.4	2.0	1.9	2.2	2.2	2.1 ^M	2.0	1.9	2.1	2.0	2.0	2.0	2.1	2.1	(2.4) ²	2.3 ^F	2.0
5	1.9	1.9 ⁵	1.9	1.9	2.0	2.2	2.2	2.2	G	G	1.9	1.9 ⁵	2.2	1.9	2.0	2.0	2.1	2.0	A	J ^A	2.3	2.2	A	A	A
6	2.3 ^F	(2.2) ^F	2.1 ^F	2.2	2.3 ^F	2.3 ^F	2.3 ^F	2.3 ^F	2.4	1.9	2.3	2.3	2.1	2.1	1.8	1.7	2.2	2.1	2.2	2.1	2.2	2.2	2.2	2.2	2.1
7	2.0 ⁵	2.0	2.0	2.1	2.1 ⁵	2.3	2.4	2.4	2.2	2.3	(2.3) ^F	2.1	2.0	2.1	1.9	2.0	2.1	2.2	2.2	2.2	2.2	2.3	2.2	(2.2) ³	2.2
8	2.1	2.1 ^F	2.2 ^F	2.2	2.1	2.3	2.3	2.3	2.3 ^M	2.1 ^M	2.1 ^M	1.8	2.2	2.0	2.1	1.8	1.9	2.3	2.1	2.0	2.2	2.1	2.1	2.2	2.1
9	2.1 ⁵	2.3 ^F	2.1	(2.1) ⁵	2.2	2.2	2.1 ^M	2.1 ^M	1.4 ^M	2.2	G	G	G	1.8 ^K	1.8 ^K	1.7 ^K	1.9 ^K	1.7 ^K	2.0 ^K	2.2 ^K	2.2 ^K	2.3 ^K	2.3 ^K	2.0 ^K	2.0 ^K
10	(2.0) ⁵	(2.0) ⁵	(2.2) ⁵	(2.1) ⁵	2.1	2.4	2.3	2.3	2.4	2.2	2.2	2.2	2.2	2.2	2.1	2.1	2.1	2.0	2.0	2.1	2.2	(2.2) ³	2.1	2.0	2.0
11	1.9	1.9	2.0	2.0	2.1	2.2	2.4	2.4	2.2	2.0	2.0	1.8	2.0	2.0	1.7	1.9	1.9	2.1	2.1	2.1	2.1	2.2	2.2	2.2	2.0
12	1.9	1.9	A	2.1	A	(2.1) ⁵	A	2.1	2.4	2.3	2.3 ^M	2.4 ^M	1.7	2.1	2.0	2.0	2.0	2.1	A	2.2	2.1	2.1	2.2	2.3	2.2 ^F
13	2.1	A	2.0	2.2 ^F	2.1 ⁵	A	A	A	G	A	A ^K	2.1 ^M	1.9	1.8	A	1.8	A	2.0	2.0	A	2.2	2.2	2.2	A	A
14	2.0 ⁵	2.1 ^F	2.2 ^F	2.2	A	A	A	2.0 ^M	2.4 ^M	2.1	2.0	2.1 ^M	2.1	2.0	1.7	1.9	1.7	2.0 ^K	A ^K	A ^K	A ^K	2.2 ^K	2.1 ^F	2.2 ^K	2.0 ^K
15	2.1 ^K	2.2 ^K	2.1	2.2 ^F	(2.2) ⁵	2.3	2.3	2.3	2.3	2.1	(1.7) ⁵	2.1	2.2	2.0	2.1 ^M	2.2	2.0	2.1	2.3	2.1	2.0	2.2	2.1	2.0	2.3
16	2.2	2.0	2.0	2.0	2.2 ^F	2.2	2.2	2.2	2.3	2.3	2.3	2.3	2.2	2.0	2.1 ^M	2.2	2.0	2.0	2.1	2.2	2.3	2.3	A	2.1	A
17	2.3	2.1	A	2.2	2.0	2.2	2.2	A	2.5	2.1	2.1	2.0 ^M	2.2	2.1	2.0	2.1	2.2	2.2	2.2	2.2	2.3	(2.3) ⁵	2.4	2.2 ⁵	2.0
18	2.0	2.1	2.0	A	2.1 ^F	2.2	2.2	2.2	T	A	2.2	2.0	1.9 ^M	1.1	1.8	1.7	2.0	2.0	2.0	2.0	A	A	2.5	(2.3) ⁵	2.3 ⁵
19	A	A	2.0	A	2.1	2.4 ^F	2.1	2.1	2.1	1.9	A	(1.8) ⁵	2.0	1.6	1.7	2.3 ^M	2.1	2.1	2.1	2.1	2.1	(2.2) ⁵	2.1	2.2 ⁵	2.2
20	2.1	2.0 ⁵	(2.2) ⁵	(2.2) ⁵	(2.2) ⁵	2.3	2.3	2.0	1.8	1.9	(1.1) ⁵	2.1	1.7	1.5	1.7	1.8	1.7	2.0	2.1	2.2	(2.3) ⁵	2.2	2.2	(2.2) ⁵	(2.1) ⁵
21	(2.0) ⁵	(2.1) ⁵	(2.1) ⁵	5	5	2.2	1.7	1.7	1.8	2.5	(1.2) ⁵	5	1.1	2.1	2.2	1.9 ^M	1.8	2.0	A	2.1	2.3	2.4	2.5	2.0	2.0
22	2.0	(1.9) ⁵	(2.1) ⁵	2.0	2.1	2.5	2.4	2.4	2.0	2.0	1.8	1.6	1.8	1.4 ^M	1.7 ^M	2.2	1.7	2.1	2.1	2.0	(2.1) ⁵	(2.3) ⁵	2.2	2.2	2.1
23	2.1	2.1	(2.0) ⁵	J ⁵	(2.1) ⁵	2.5	2.5	1.4 ^M	2.3	2.4	2.3	A	A	5	2.0	A	A	2.0	M	M	M	M	2.2	2.2	2.1
24	2.2	A	(2.1) ⁵	(2.1) ⁵	(2.0) ⁵	2.2	2.2	2.2	1.8	(1.4) ²	2.2 ^M	2.3	A	1.5	1.7	1.6	2.0	2.1	2.2	2.2	2.2	2.1	A	(2.1) ⁵	
25	(2.2) ⁵	2.2 ⁵	2.2	2.3 ^F	2.3	2.5	2.5	2.4	2.3	2.1 ⁵	2.3 ^M	2.3	A	1.4	2.1	2.0	2.0	1.1	2.2	2.2	2.2	2.3	2.3	2.1	
26	2.2	A	A	A	A	2.1	A	2.1	A	A	A	(2.1) ⁵	2.1	1.7	2.2	A	A	A	2.3	A	(2.3) ⁵	(2.3) ⁵	2.2	2.2	2.1
27	2.0 ⁵	2.1 ^F	(2.2) ⁵	(2.2) ⁵	2.0 ^F	2.5	2.5	2.5 ^M	1.7	2.2 ^M	A	A	2.1	2.2	1.7	2.0	1.7 ⁵	2.0	A	A	2.2	2.3	2.3	2.2	2.2 ⁵
28	(2.2) ⁵	2.2 ⁵	(2.2) ⁵	A	A	2.2	2.5	2.5	2.2	2.3 ^M	2.1 ⁵	2.4	A	2.2	2.1	1.9	2.0	2.0	2.0	2.1	(2.1) ^A	2.2	2.3 ⁵	2.2	2.2 ⁵
29	2.2 ^F	2.2 ⁵	2.2	2.1 ^F	2.4 ⁵	(2.2) ⁵	1.8	1.8	1.8	1.4	2.2	2.1	2.0	1.8	1.6	1.7	1.7	1.7	2.0	2.1	2.2	2.3	2.3	2.2	2.2
30	2.1	2.1	2.1	(2.0) ⁵	2.0	2.2	2.2	2.2	2.1	2.1	1.9 ^M	2.2	2.2	2.0	2.2	1.8	2.0	2.2	2.1	2.1	2.2 ⁵	2.2	2.3	2.2	2.1
31	2.0	2.2	2.2 ^F	2.3 ^F	J ⁵	2.2 ^M	1.7 ^M	1.7 ^M	2.1	2.4	2.2	2.0	2.1	2.0	2.0	J ⁵	1.8	2.0	2.1	M	2.2	2.3	2.2	2.2	2.2
Median	2.1	2.1	2.1	2.2	2.1	2.2	2.2	2.2	2.2	2.1	2.1	2.1	2.2	2.0	2.0	1.7	2.0	2.0	2.1	2.1	2.2	2.2	2.2	2.1	2.1
Count	30	27	27	25	26	28	28	28	28	28	26	27	27	29	29	28	28	29	24	24	27	29	28	28	28

Sweep 1.0 — Mc to 25.0 Mc in 0.25 min

Manual ☐ Automatic ☒

TABLE 82

Control Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D. C.

(M3000)F2
(Characteristic)
Observed at Washington, D.C.
May 1954
(Month)

Form adopted June 1946

National Bureau of Standards
(Institution)
Scaled by: E.J.W., J.W.P., J.J.S.

Calculated by: E.J.W., J.W.P., J.J.S.

Lat. 38.7°N, Long. 77.1°W

Mean Time
75°W

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	(1.9) ^F	(2.8) ^F	(3.0) ^F	(3.3) ^F	3.2	3.3	3.4	2.8	(2.8) ^S	3.2	3.4	3.0	3.2	3.0	3.1	3.1	3.1	3.1	3.3	3.3	3.2	3.2	3.0	(3.0) ^S
2	2.9	3.0	3.2	3.2	3.2	3.3	3.3	3.4	3.0	2.9	3.2	2.7 ^H	3.2	3.1	3.2	3.1	3.1	3.1	3.2	3.3	3.2	3.2	3.1	3.1
3	3.1	2.9	3.1	3.2	3.2	3.3	3.3	3.5	3.4	3.1	3.3 ^H	3.2	3.4	3.2	3.1	3.2	3.3	3.3	3.3	3.3	3.3	3.4	3.4	3.2 ^F
4	3.3 ^F	3.2 ^F	3.2 ^F	3.3 ^F	3.4 ^F	3.3 ^F	3.4 ^H	3.5	3.0	3.1 ^H	3.3	3.3	3.1 ^H	3.0	2.9	3.1	3.0	3.0	2.9	3.1	3.1	(3.4) ^S	3.3 ^F	3.0
5	2.9	2.9	A	(2.9) ^S	2.9	3.0	3.3	G	G	2.8	2.9 ^S	3.3	2.8	S	2.9	3.1	A	A	J	A	3.3	3.2	A	A
6	3.3 ^F	(3.3) ^F	3.1 ^F	3.2	3.3 ^F	3.4 ^S	3.4	3.4	2.9	3.4	C	3.1	3.1	2.7	2.8	3.2	3.1	3.2	3.1	3.2	3.2	3.2	3.2	3.1
7	3.0 ^F	3.0	3.0	3.1	3.1 ^F	3.3	3.4	3.2	3.3	(3.3) ^S	3.1	3.0	3.1	2.9	3.0	3.1	3.3	3.2	3.3	3.2	3.2	(3.2) ^S	3.2	
8	3.2	3.2 ^F	3.2 ^F	3.1	3.1	3.4	3.4	3.4 ^H	3.1 ^H	3.1 ^H	2.8	3.2	3.0	3.1	2.7	2.9	3.0	3.1	3.0	3.2	3.1	3.1	3.2	3.2
9	3.1 ^F	3.3 ^F	3.1	(3.2) ^S	(3.0) ^F	3.2	3.1 ^H	2.9 ^H	3.2	G	G	2.7 ^K	2.7 ^K	2.7 ^K	2.6 ^K	2.9 ^K	2.9 ^K	3.0 ^K	3.2 ^K	3.2 ^K	3.3 ^K	3.3 ^K	3.0 ^K	3.0 ^K
10	(3.0) ^F	(3.0) ^F	(3.2) ^S	(3.1) ^S	3.1 ^F	3.4	3.3	3.5	3.2	3.2	3.3	3.2	3.2	3.1	3.1 ^F	3.2	3.0	3.0	3.1	3.2	(3.2) ^S	3.1	3.0	3.0
11	2.9	2.9	3.0	3.0	3.1	3.3	3.4	3.3	3.0	3.0	2.8	3.0	3.0	2.9	2.9	2.9	3.2	3.1	3.1	3.1	3.3	3.3	3.3	3.0
12	2.9	2.9	A	3.1	A	(3.0) ^S	3.1	3.4	3.4	3.3 ^H	3.4 ^H	2.6	3.1	3.0	3.0	3.0	3.1	A	3.2	3.1	3.1	3.2	3.4	3.3 ^F
13	3.1 ^F	A	3.0	3.2 ^F	3.1 ^F	A	A	A	2.9	A	3.1 ^H	2.9	2.7	A	2.8	A	3.0	3.0	A	3.3	3.3	3.2	A	A
14	3.0 ^F	3.1 ^F	3.1 ^F	3.2 ^F	3.2	A	A	A	A	A	A	G	2.2 ^K	G	2.4 ^K	2.9 ^K	3.0 ^K	A	A	A	3.3 ^K	3.1 ^K	3.2 ^K	3.0 ^K
15	3.1 ^K	3.3 ^F	3.1	3.2 ^F	(3.3) ^S	3.4	3.0 ^H	3.4 ^H	3.1	3.0	3.1 ^H	3.1	3.0	2.9	2.8	2.9	3.1	3.3	3.1	2.9	3.3	3.1	3.0	3.3
16	3.2	3.0	2.9	3.1	3.2 ^F	3.2	G	3.3	2.9	(2.6) ^S	3.2	3.2	3.0	3.0 ^H	3.2	3.0	3.0	3.1	3.2	3.3	3.4	A	3.1	A
17	3.2	3.1	A	3.3	3.0	3.2	A	3.5	3.1	3.1	3.0 ^H	3.2	3.1	3.0	3.1	3.2	3.2	3.2	3.2	3.3	(3.3) ^S	3.4	3.2 ^S	3.0
18	3.0	3.1 ^F	3.0	A	3.1 ^F	3.2	3.2	T	A	3.2	3.0	2.9 ^H	2.6	2.7	2.8	3.0	2.9	3.0	3.0	A	A	3.5	(3.5) ^S	3.3 ^F
19	A	A	3.0	A	3.1	3.5 ^F	3.1	3.1	2.9	A	(2.8) ^S	3.0	2.4	2.8	3.0 ^H	3.1	3.1	3.1	3.1	3.1	(3.3) ^S	3.2	3.2 ^S	3.2
20	3.1	3.0 ^S	(3.2) ^S	(3.2) ^S	(3.0) ^F	3.3	3.0	2.7	2.9	(2.5) ^S	3.1	2.8	G	2.9	2.7	2.8	3.0	3.1	3.2	(3.3) ^S	3.2	3.3	(3.2) ^S	(3.0) ^S
21	(3.0) ^F	(3.0) ^F	(3.0) ^F	S	S	3.2	1.6	2.7	3.3	(2.3) ^S	G	2.9	3.1	3.2	2.6 ^H	2.8	3.0	A	3.1	3.3	3.4	3.5	3.0	2.0
22	3.0	(2.9) ^S	(3.0) ^S	3.0	3.1	3.5	3.3	3.0	3.0	2.8	2.5	2.8	2.8 ^H	2.9 ^H	3.2	2.9	3.1	3.1	3.0	(3.0) ^S	(3.4) ^S	3.2	3.3	3.1
23	3.1	3.1	(3.0) ^S	J	(3.0) ^S	3.3	3.4	3.4	3.5	3.4	A	A	G	3.0	A	A	3.0	M	M	M	M	3.2	3.2	3.1
24	3.2	A	(3.0) ^S	(3.0) ^S	(3.0) ^S	3.2	2.8 ^H	2.7	(2.8) ^S	3.2 ^H	3.3	A	2.2	2.7	2.4	3.0	3.1	3.2	3.3	3.3	3.1	A	(3.0) ^F	
25	(3.2) ^S	3.2 ^S	3.2	3.3 ^F	3.3	3.6	3.5	3.3	3.1 ^S	3.3 ^H	3.3	A	2.9	3.1	3.0	3.0	2.9	3.2	A	3.2	3.3	3.3	3.3	3.1
26	3.2	A	A	A	A	A	3.2	A	A	A	(3.0) ^S	3.1	2.9	3.2	A	A	A	A	3.3	A	(3.4) ^S	(3.3) ^S	3.2	3.1
27	3.0 ^F	3.2 ^F	(3.0) ^F	(3.2) ^F	3.0 ^F	3.5	3.5 ^H	2.6	3.2 ^H	A	A	3.1	3.2	2.9	3.0	2.9 ^S	2.9	A	A	3.3	3.4	3.3	3.3	3.2 ^S
28	(3.2) ^F	3.3 ^F	(3.3) ^F	A	A	3.3	3.6	3.3	3.3 ^H	3.1 ^S	3.2	A	3.2	3.1	2.8	3.0	3.0	3.0	3.1	(3.0) ^A	3.2	3.3 ^S	3.3	3.3 ^S
29	3.3 ^F	3.2 ^S	3.2	3.1 ^F	3.4 ^S	(3.2) ^S	2.7	2.7	2.9	3.2	3.1	2.9	2.5	2.6	2.9	2.8	3.0	3.1	3.2	3.2	3.2	3.3	3.2	3.2
30	3.1	3.1	3.1	(3.0) ^S	3.0	3.3	3.2	3.2	3.1	2.8 ^H	3.2	3.2	3.0	3.3	2.7	3.0	3.2	3.1	3.1	3.2 ^S	3.2	3.4	3.2	3.1
31	3.0	3.2	3.2 ^F	3.4 ^H	J	3.2	2.0 ^H	3.1	3.5	3.4	3.0	3.1	3.0	3.0	J ^S	2.7	3.0	3.1	M	3.3	3.4	3.2	3.2	3.2
Median	3.1	3.1	3.1	3.2	3.1	3.3	3.2	3.2	3.1	3.1	3.1	3.0	3.0	3.0	2.9	3.0	3.0	3.1	3.1	3.2	3.2	2.9	3.2	3.1
Count	30	27	27	25	26	28	28	28	26	26	27	27	31	29	26	26	29	24	24	27	29	29	26	26

Sweep 1.0 Mc to 25.0 Mc in 0.25 min

Manual ☐ Automatic ☒

TABLE 83

(M3000)FI
(Characteristic)
Observed at Washington, D.C.
(Unit)
Moy 54
(Month)

IONOSPHERIC DATA

National Bureau of Standards

Scaled by E.J.W., J.W.P., J.J.S.

(Institution)

Lat 38.7°N, Long 77.1°W

75°W Mean Time

Calculated by E.J.W., J.W.P., J.J.S.

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1							L	36	38	37H	39	42H	38	37	37	37	36H	A	A					
2							L	L	37H	38	40	39H	38	39H	39	(39)S	37	35H	36					
3							L	36	36H	37H	41H	40H	40H	38	37H	36	37	37	L	A				
4							L	37	35	A	39	40H	39H	39	38H	35	35H	35	35					
5							L	35H	37H	38H	42	40	41	38H	A	36	A	A	A					
6							36H	37H	38	36	C	39	40H	38	38	37	37	A	37					
7							A	38	37H	38	40H	38	39	38	38	37H	36H	37	L					
8							L	36H	38H	36H	38	38H	39	(39)S	37	36H	36	36	36					
9							33S	37H	38H	39H	39H	39H	40H	40H	37H	36H	36H	36H	A	X				
10							L	36H	37H	37	38H	A	(42)P	40H	37H	36	A	A	A					
11							L	38	36H	38	39H	39	37	37	35	A	A	A	A					
12							34	37	37	38	37H	38H	A	A	A	A	A	A	A					
13							A	A	39S	A	40	40H	39H	A	37H	A	A	A	A					
14							A	38	A	A	A	40H	39H	39H	38H	38H	A	A	A	X				
15							36H	(38)S	37H	(36)S	37H	42	39	38	37	37	38	35H	A					
16							34	35H	37H	39	40	38H	(39)S	38H	39H	36	36	36H	A					
17							36	40	A	39	41H	40	39	38	38	A	37H	37H	L					
18							35	T	A	39	40	39	39	(40)S	39H	37H	36	35	36					
19							36	36H	37	A	A	39	(40)S	39	39H	39	37H	35H	36					
20							35	37	38	41	40	36H	39	39H	38	36H	37	36	A					
21							35	39	38	39H	40H	41	41	41	39	38	37	37	A					
22							(38)S	38H	39	39H	42	39H	39	39	39	A	35	36H	A					
23							35	37H	39H	40H	A	A	41H	A	A	A	A	M	M					
24							38	40	38	40H	40H	A	41H	38	37	36H	A	35	36H					
25							A	(37)S	37H	40H	41	A	39	39	38	38	36	A	A					
26							38	A	A	A	41H	41H	(42)S	A	A	A	A	A	A					
27							L	40	38	A	A	A	38H	38	40	40	A	A	A					
28							L	36	37H	38	A	A	40H	40	37H	38	38	36	L					
29							34H	36	38H	39	40	39H	39S	38H	38H	37	36	35	36					
30							36	37	37	38H	39H	41H	37H	38H	38	37	37H	36	A					
31							35H	38	A	41H	(42)P	41H	40H	39H	39H	38	37	37	M					
Median							35	37	38	38	40	39	39	39	38	37	36	36	36					
Count							16	17	17	17	17	17	17	17	17	17	17	17	17					

Sweep 1.0 Mc to 25.0 Mc in 0.25 min
Manual ☐ Automatic ☒

TABLE 84

Central Radio Propagation Laboratory, National Bureau of Standards, Washington 25, D.C.

(M1500)E

(Unit)

May 1954

(Month)

Observed at Washington, D.C.

Lat 38.7°N, Long 77.1°W

IONOSPHERIC DATA

Form adopted June 1948

National Bureau of Standards

(Institution)

Scaled by: E.J.W., J.W.P., J.J.S.

Calculated by: E.J.W., J.W.P., J.J.S.

Day	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1							4.3	4.3	4.3	4.3	4.4	(4.2)P	A	4.4H	A	A	B	4.0	S					
2							4.2H	4.5H	4.3	4.4	4.5	A	4.3H	4.4	A	(4.4)A	(4.3)P	4.5	4.5					
3							4.4H	4.4H	4.4	4.3	4.4H	4.4H	4.5	4.5H	(4.5)S	A	(4.4)S	4.5	S					
4							4.4H	4.5	4.5H	4.4	4.5	4.5	A	4.5	4.5	4.3H	4.4H	4.1H	4.3					
5							4.4	4.3H	4.3H	4.3	(4.4)A	A	A	4.4	4.4	(4.4)B	4.2H	4.3	(4.4)P					
6							A	A	(4.3)A	4.4	C	(4.4)A	A	4.3	4.2	(4.2)P	4.1	4.2	A					
7							A	4.2	4.4	4.5	4.4	(4.5)A	A	A	S	A	4.3	4.3	4.3					
8							4.5H	4.5H	A	A	4.5	4.4	4.4H	4.5	4.4H	4.4	4.4H	4.3	4.2					
9							A	4.4	4.4H	A	4.4	4.4H	(4.5)P	4.5H	4.4H	4.3H	4.4H	4.4H	4.4H					
10							4.5	(4.4)P	4.4	4.5	(4.5)A	A	A	A	4.3	4.3	4.3	4.2	(4.2)A					
11							S	4.3	4.5S	4.5	4.4	4.4	4.3H	4.3H	4.3	4.4	4.4	4.4	4.4					
12							4.4	4.3	4.4H	4.3H	4.3H	4.5H	4.5H	4.4	4.4	(4.5)A	4.5	4.3	4.5					
13							4.5	4.3H	4.5	4.5	4.4	A	A	A	4.4H	4.4	4.4	4.4	4.4					
14							(4.5)A	4.5H	4.5	AH	AH	AH	AH	AH	AH	4.4H	4.4H	4.4H	4.4H					
15							A	A	4.4	4.4H	4.5	4.5H	4.4	4.5	(4.6)A	(4.5)A	4.3	4.4	4.3H					
16							4.5	4.5	4.6	4.6	4.5	4.4	4.5	4.5H	(4.5)P	A	A	A	A					
17							4.4H	4.5H	4.4	4.4	4.4H	4.5H	A	A	A	A	4.4	4.5	4.6	(4.5)P				
18							4.4	T	4.5	4.5	4.6	4.5	A	(4.5)A	A	4.4	(4.4)B	4.3	(4.3)P					
19							4.4	4.4	4.5	4.4H	4.5	4.5	4.5	4.5	(4.5)P	4.5	A	4.4	4.4					
20							A	A	4.4H	4.4H	(4.4)P	(4.5)B	4.3H	(4.4)P	(4.4)P	4.4	(4.2)B	4.3H	4.4					
21							(4.4)A	4.4H	4.5	(4.5)P	A	A	(4.5)P	B	4.4	4.5	4.4	4.3	A					
22							S	4.4H	4.5	4.5H	(4.6)A	4.6	(4.5)P	A	(4.6)P	4.4	(4.4)A	(4.4)S	(4.4)H					
23							4.4H	4.4H	(4.4)A	4.5H	4.5H	4.5	(4.4)P	B	4.4H	4.4	M	M	M					
24							(4.5)H	A	A	(4.5)P	4.5	4.5	(4.4)P	(4.4)P	S	4.4H	4.4	4.4	4.6					
25							4.5	4.6	4.5	A	A	A	A	A	A	4.5	4.4	4.5	4.5H					
26							4.4H	4.4	4.5	4.5	4.5	4.6	4.4	4.5	A	4.4H	4.4	4.5	4.6					
27							4.5	4.5	4.4	4.6	4.6	(4.5)A	(4.5)A	A	4.4H	4.4	(4.4)P	4.4	4.5					
28							4.6H	4.5	4.5S	4.5H	A	A	A	4.4	4.4	4.4	4.5H	4.5	4.6					
29							4.5H	4.4	4.4	A	A	A	4.5	4.5	A	A	A	(4.4)P	4.5					
30							4.4	4.4	4.4H	4.5	4.5	A	A	A	A	4.4H	4.5H	4.5H	4.5H					
31							4.5H	4.4	4.4	4.5	4.5	4.5H	4.3H	(4.4)A	4.4	4.5	4.4	4.4H	M					
Median							4.4	4.4	4.4	4.5	4.5	4.5	4.4	4.4	4.4	4.4	4.4	4.4	4.4					
Count							24	26	29	26	25	21	18	20	21	26	26	29	24					

Sweep 1.0—Mc to 25.0 Mc in 0.25 min

Manual ☐ Automatic ☒

Table 85

Ionospheric Storminess at Washington, D. C.May 1954

Day	Ionospheric character*		Principal storms		Geomagnetic character**	
	00-12 GCT	12-24 GCT	Beginning GCT	End GCT	00-12 GCT	12-24 GCT
1	2	2			2	1
2	1	1			3	2
3	1	1			2	1
4	1	2			3	3
5	3	2			2	2
6	2	2			3	1
7	1	1			1	2
8	0	0			2	3
9	0	4	1100	----	3	2
10	2	2	----	2400	2	3
11	3	1			4	3
12	3	1			2	2
13	2	3			2	3
14	2	5	0900	----	3	2
15	1	2	----	0100	2	3
16	1	2			2	1
17	3	2			1	1
18	2	3			3	3
19	3	2			3	3
20	1	3			2	3
21	1	2			3	3
22	3	2			2	2
23	1	3			2	1
24	2	3			2	2
25	1	1			2	1
26	2	2			2	2
27	2	2			2	1
28	2	1			1	2
29	1	3			3	2
30	2	1			2	1
31	1	2			1	2

*Ionosphere character figure (I-figure) for ionospheric storminess at Washington, D. C., during 12-hour period, on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

**Average for 12 hours of Cheltenham, Maryland, geomagnetic K-figures on an arbitrary scale of 0 to 9, 9 representing the greatest disturbance.

----Dashes indicate continuing storm.

Radio Propagation Quality Figures
(Including Comparisons with Short-Term and Advance Forecasts)

April 1954

Day	North Pacific 9- hourly quality figures			Short-term fore- casts issued at:			Whole day quality index	Advance forecasts (Jp reports) for whole day; issued in advance by:		
	03 to 12	09 to 18	18 to 03	02	09	18		1-4 days	4-7 days	8-25 days
1	7	6	6	6	6	6	7	5	6	
2	5	5	7	6	5	6	6	6	6	
3	6	6	7	6	5	7	7	6	6	
4	6	6	7	6	5	6	6	6	6	
5	6	6	7	6	5	7	7	5	5	
6	5	6	6	7	6	7	6	5	5	
7	5	6	7	6	6	7	6	6	6	
8	6	6	7	6	6	7	6	6	6	
9	6	6	7	6	6	6	6	6	5	
10	5	6	7	6	5	6	6	(4)	(4)	X
11	6	6	5	6	5	7	6	(4)	(4)	X
12	(4)	6	5	5	(4)	6	5	(4)	(4)	X
13	5	(4)	6	(4)	(4)	6	5	5	5	
14	(4)	5	5	5	5	6	5	5	5	
15	5	6	5	6	5	6	5	5	5	
16	5	5	6	5	(4)	6	5	6	6	
17	6	6	6	5	5	6	6	5	6	
18	5	6	7	5	5	6	6	5	6	
19	6	5	6	6	5	6	6	(4)	(4)	X
20	5	5	6	5	5	5	6	(4)	(4)	X
21	5	(4)	6	6	5	6	5	5	5	
22	5	5	7	6	5	7	6	6	5	
23	6	6	6	6	5	7	6	6	6	
24	6	6	6	6	5	6	6	6	6	
25	5	5	7	6	5	7	5	6	6	
26	6	6	6	6	6	7	6	6	6	
27	5	6	6	5	5	6	6	6	5	
28	5	6	5	5	6	6	6	5	6	
29	5	5	6	5	6	7	5	6	6	
30	5	5	6	6	5	6	5	6	6	

Score:

Quiet Periods	P	16	14	15	14	14
	S	11	13	14	10	11
	U	1	0	1	2	1
	F	0	1	0	4	4

Disturbed Periods	P	0	1	0	0	0
	S	2	1	0	0	0
	U	0	0	0	0	0
	F	0	0	0	0	0

Scales:

Q-scale of Radio Propagation Quality

- (1) - useless
- (2) - very poor
- (3) - poor
- (4) - poor to fair
- 5 - fair
- 6 - fair to good
- 7 - good
- 8 - very good
- 9 - excellent

K-scale of Geomagnetic Activity

0 to 9, 9 representing the greatest disturbance; $K_{ch} \geq 4$ indicates significant disturbance, enclosed in () for emphasis

Scoring: (beginning October 1952)

- P - Perfect: forecast quality equal to observed
- S - Satisfactory: (beginning October 1952) forecast quality one grade different from observed
- U - Unsatisfactory: forecast quality two or more grades different from observed when both forecast and observed were ≥ 5 , or both ≤ 5
- F - Failure: other times when forecast quality two or more grades different from observed

Symbols:

X - probable disturbed date

Note: All times are UT (Universal Time or GCT)

Radio Propagation Quality Figures
(Including Comparisons with Short-Term and Advance Forecasts)

April 1954

Day	North Atlantic 6-hourly quality figures				Short-term forecasts issued about one hour in advance of:				Whole day quality index	Advance forecasts (J-reports) for whole day; issued in advance by:			Geomag- netic K _{Ch}	
	00 to 06	06 to 12	12 to 18	18 to 24	00	06	12	18		1-4 days	4-7 days	8-25 days	Half day (1) (2)	
1	6	(4)	7	7	6	5	6	7	6	6	6		2	2
2	6	5	7	6	6	5	7	7	6	6	6		2	3
3	5	(4)	7	6	5	5	7	6	6	6	6		3	3
4	(4)	(4)	7	7	6	(4)	7	6	6	6	6		3	3
5	6	5	7	7	5	5	7	7	6	6	6		3	2
6	6	5	7	7	6	6	7	7	6	6	6		2	2
7	6	5	7	7	7	6	7	7	6	6	5		2	1
8	6	5	7	6	6	5	7	7	6	6	5		1	3
9	6	5	7	6	6	5	6	6	6	6	6		2	2
10	6	(4)	7	7	6	(4)	6	6	6	5	5		3	2
11	6	6	7	6	6	5	7	7	7	(4)	(4)	X	3	(4)
12	(3)	(3)	5	5	5	(2)	5	(4)	(4)	(4)	(4)	X	(5)	(4)
13	(2)	(3)	6	6	(3)	(2)	5	5	(4)	(5)	5		2	(4)
14	(4)	(4)	6	6	(4)	(3)	6	6	5	(4)	6		3	2
15	5	(4)	7	6	5	(4)	6	5	5	(4)	6		3	3
16	5	(4)	6	6	5	5	6	6	5	5	5		2	2
17	6	(4)	7	7	5	(4)	6	6	6	6	5		2	2
18	5	5	7	7	6	5	7	7	6	6	5		3	2
19	6	(4)	6	6	5	5	6	6	5	(4)	(4)	X	3	3
20	6	(4)	7	6	5	(4)	6	6	6	(4)	(4)	X	(4)	3
21	5	(4)	7	6	6	(4)	7	7	6	(4)	(4)	X	3	2
22	5	5	7	6	5	(4)	6	6	6	5	5		2	2
23	6	5	7	6	6	5	7	7	6	6	6		3	3
24	5	5	6	6	5	(3)	6	6	6	6	6		3	3
25	5	6	7	7	5	(4)	6	5	6	6	6		3	2
26	5	5	7	6	5	5	3	6	6	6	6		3	3
27	(4)	(3)	6	6	6	5	5	6	(4)	6	6		(4)	2
28	6	5	7	7	5	5	7	6	6	6	6		2	2
29	6	5	7	7	5	5	3	6	7	6	6		2	2
30	6	5	7	6	6	5	6	6	6	6	6		3	3
Score:														
Quiet periods					P	16	10	17	16		13	14		
					S	9	4	13	13		6	10		
					U	0	1	0	1		0	0		
					F	0	1	0	0		3	3		
Disturbed periods					P	1	6	0	0		1	1		
					S	1	7	0	0		1	1		
					U	1	1	0	0		0	0		
					F	2	0	0	0		1	1		

Scales:

Q-scale of Radio Propagation Quality

- (1) - useless
- (2) - very poor
- (3) - poor
- (4) - poor to fair
- 5 - fair
- 6 - fair to good
- 7 - good
- 8 - very good
- 9 - excellent

K-scale of Geomagnetic Activity

0 to 9, 9 representing the greatest disturbance; K_{Ch} ≥ 4 indicates significant disturbance, enclosed in () for emphasis

Scoring: (beginning October 1952)

- P - Perfect: forecast quality equal to observed
- S - Satisfactory: (beginning October 1952) forecast quality one grade different from observed
- U - Unsatisfactory: forecast quality two or more grades different from observed when both forecast and observed were 2, 3, or both 4, 5
- F - Failure: other times when forecast quality two or more grades different from observed

Symbols:

X - probable disturbed date

Note: All times are UT (Universal Time or GCT)


Short-Term Forecasts---April 1954

— forecast

• individual reports of quality

x CRPL observation (not in consensus)

(adjusted to CRPL scale)



Outcome of Advance Forecasts (1 to 4 days ahead) --- April 1954




Table 88a

Coronal observations at Climax, Colorado (5303A), east limb

Date (UT)	Degrees north of the solar equator																	Degrees south of the solar equator														
	50	45	40	35	30	25	20	15	10	5	0	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90			
1954																																
May 2.9	-	-	-	-	-	-	-	-	-	-	1	4	5	4	3	1	1	1	1	1	1	1	-	-	-	-	-	-	X	X		
3.7	-	-	-	-	-	-	-	-	-	-	2	3	5	4	2	1	-	-	1	1	1	-	-	-	-	-	-	-	-	-		
5.6	-	-	-	-	-	-	-	-	-	-	1	1	2	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
6.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
7.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
8.7a	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X	X		
9.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
11.7a	X	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
12.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
14.6a	-	-	-	-	-	-	-	-	-	-	1	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
17.6	-	-	-	-	-	-	-	-	-	-	1	1	1	-	-	-	1	1	1	1	1	1	1	-	-	-	-	-	-	-		
18.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	3	3	3	-	-	-	-	-	-	-	-	-		
20.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
21.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
24.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25.6	-	-	-	-	-	-	-	-	-	-	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
26.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
27.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
28.6	-	-	-	-	-	-	-	-	-	-	1	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
30.7a	-	-	-	-	-	-	-	-	-	-	-	2	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
31.7	-	-	-	-	-	-	2	2	-	-	-	-	-	-	-	-	1	1	1	-	-	-	-	-	-	-	-	-	-	-		

Table 89a

Coronal observations at Climax, Colorado (6374A), east limb

Date (UT)	Degrees north of the solar equator																	Degrees south of the solar equator																	
	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
1954																																			
May 2.9	2	2	2	2	1	1	1	1	1	2	3	3	4	5	5	4	3	3	2	2	2	3	5	4	3	3	2	2	2	2	2	2	X	X	
3.7	3	3	2	2	1	1	1	1	1	1	2	2	3	4	5	5	5	4	5	5	3	6	3	5	6	5	3	2	1	1	1	1	2	3	2
5.6	2	3	2	2	1	1	1	2	2	1	1	3	3	2	6	5	5	6	5	5	6	6	4	3	2	2	2	1	1	1	1	1	2	2	
6.6	2	2	1	1	1	1	1	1	1	1	2	3	4	5	4	4	4	5	5	3	3	2	2	2	1	1	1	1	1	1	1	1	1		
7.6	1	2	1	1	1	1	1	1	1	1	3	3	3	3	4	5	4	4	4	3	3	3	3	2	2	2	1	1	1	1	1	1	1	2	
8.7a	X	2	1	1	1	-	-	-	-	-	1	1	1	3	3	3	3	2	1	1	1	1	2	1	1	1	1	-	-	-	-	-	X	X	
9.7a	3	2	1	1	1	1	1	1	1	1	3	3	2	3	4	4	3	3	3	3	3	3	3	3	3	1	1	1	1	1	1	2	2		
11.7a	X	X	-	-	-	-	-	-	-	-	3	3	3	3	3	5	6	5	3	3	3	3	3	3	3	3	2	2	2	2	2	2	2	2	
12.7	2	2	1	1	1	1	1	1	-	1	2	2	2	2	2	3	4	3	4	4	3	3	4	4	3	2	1	1	1	1	1	1	2	2	
14.6a	2	1	1	1	1	1	1	1	1	1	2	2	2	3	3	3	3	3	3	3	2	2	2	2	1	1	1	1	1	1	1	1	2	2	
17.6	2	2	2	1	1	1	1	1	2	2	3	3	3	4	4	5	6	6	6	6	6	5	5	5	3	2	1	1	1	1	1	2	2		
18.7	2	1	1	1	1	1	1	1	1	2	3	5	5	4	4	5	6	6	6	5	5	5	5	5	3	1	1	1	1	1	1	1	2	2	
20.7a	2	2	2	2	-	-	-	-	1	1	1	2	2	3	4	4	5	5	5	2	3	3	3	4	3	2	1	1	1	1	1	1	2	2	
21.6a	2	2	1	1	1	1	1	1	1	1	2	3	3	3	3	6	5	3	3	3	3	3	2	2	2	1	1	1	1	1	1	1	2	2	
24.6	2	2	2	1	1	1	1	1	1	1	1	1	2	3	4	3	4	3	5	4	4	4	5	4	3	1	1	1	1	1	1	1	2	2	
25.6	2	2	2	1	1	1	1	1	1	1	1	1	2	2	2	3	3	3	3	4	4	4	3	2	2	1	1	1	1	1	1	1	1	2	
26.6a	2	2	-	-	-	-	-	-	-	-	1	1	2	2	2	2	3	3	3	1	1	1	1	1	2	1	-	-	-	-	-	-	-	-	
27.7a	2	2	2	1	1	-	-	-	-	-	-	1	1	1	3	3	2	1	1	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
28.6	2	2	2	2	1	1	1	1	1	1	2	4	3	3	3	2	2	2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
30.7a	3	3	2	2	2	1	1	1	1	1	1	1	1	3	4	5	4	4	3	3	4	4	3	2	2	2	1	1	1	1	1	1	2	2	
31.7	2	2	2	2	1	1	1	-	-	2	2	3	3	3	4	5	6	6	6	6	6	5	4	4	3	3	1	1	1	1	1	2	3	3	2

Table 90a

Coronal observations at Climax, Colorado (6702A), east limb

The 6702A coronal line was not visible on any of the observation dates in May at the position angles indicated for the 6374A line.

Table 88b

Coronal observations at Climax, Colorado (5303A), west limb

Date GCT	Degrees south of the solar equator																			Degrees north of the solar equator																			
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
1954																																							
May 2.9	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	
3.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
6.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
7.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.7a	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
9.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
11.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X		
12.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
14.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
17.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
18.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
21.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
24.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
26.6a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
27.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
28.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
31.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

Table 89b

Coronal observations at Climax, Colorado (6374A), west limb

Date GCT	Degrees south of the solar equator																			Degrees north of the solar equator																			
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
1954																																							
May 2.9	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	2	
3.7	2	2	2	2	2	2	1	1	1	1	1	3	4	4	4	5	4	4	5	5	5	3	4	4	6	4	3	2	2	2	2	3	3	3	3	3	3	3	
5.6	2	2	1	1	1	1	1	1	1	1	1	1	5	5	3	4	4	4	5	4	4	5	5	4	4	3	3	2	1	1	1	1	2	2	2	2	2	2	
6.6	1	2	2	2	1	1	1	1	1	1	1	2	3	4	4	3	3	3	4	3	4	3	3	2	2	2	2	2	1	1	1	1	1	1	1	2	2	2	
7.6	2	2	1	1	1	1	1	1	1	1	1	3	2	2	2	3	3	3	3	3	4	3	3	2	2	2	2	3	2	1	1	1	1	1	1	1	1	1	
8.7a	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
9.7a	2	2	2	2	1	1	1	1	1	1	1	2	3	3	4	4	4	4	4	4	3	3	3	3	3	3	2	2	1	1	1	1	1	1	1	2	3		
11.7a	2	2	2	2	1	1	1	1	2	3	3	4	4	5	5	5	5	5	5	5	5	5	5	5	5	3	2	2	2	2	2	2	2	2	2	2	X		
12.7	2	1	1	1	1	1	1	1	1	1	3	3	3	3	3	3	3	3	4	4	4	3	3	3	2	2	1	1	1	1	1	1	1	1	2	2	2		
14.6a	2	1	1	1	1	-	-	-	-	-	-	-	-	3	3	3	3	3	3	3	4	5	3	3	3	3	3	3	1	1	1	1	1	2	2	2	2		
17.6	2	2	2	2	2	2	1	1	1	1	1	1	1	4	4	4	3	5	4	5	4	2	3	3	2	2	2	1	1	1	1	1	1	2	2	2	2		
18.7	2	2	2	2	2	1	1	1	1	1	1	3	4	5	5	5	5	5	5	5	3	3	2	2	2	2	2	1	1	1	2	2	2	2	2	2	2		
20.7a	2	2	3	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	3	2	2	2	2	2	2	2	2	2	1	1	1	1	1	2	2	2	2		
21.6a	2	2	2	2	1	1	1	1	1	2	2	3	4	4	4	3	3	3	3	3	3	4	4	4	2	2	2	2	1	1	1	1	1	1	2	2	2		
24.6	2	2	2	2	2	1	1	1	1	1	1	1	3	4	3	2	2	2	2	3	3	3	3	3	3	3	3	4	5	1	1	1	1	1	1	2	2		
25.6	2	2	2	2	2	2	2	1	1	1	1	2	2	2	2	2	3	4	3	3	3	3	3	3	3	3	3	4	3	2	1	1	1	1	1	1	2		
26.6a	-	-	-	-	-	-	-	-	-	-	1	1	1	1	1	2	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
27.7a	-	-	-	-	2	2	2	2	2	2	2	2	2	2	2	2	3	3	3	3	3	3	3	3	3	3	3	2	2	1	1	1	1	1	1	2	2		
28.6	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	2	2	3	3	2	3	3	3	3	3	2	2	2	1	1	1	1	1	1	2	2			
30.7a	2	2	2	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	1	1	1	1	1	2	2	3	3		
31.7a	2	2	1	1	1	1	1	1	1	1	1	1	2	5	5	6	6	6	6	6	6	5	6	5	5	5	4	3	2	2	2	2	3	3	3	3	3		

Table 90b

Coronal observations at Climax, Colorado (6702A), west limb

The 6702A coronal line was not visible on any of the observation dates in May at the position angles indicated for the 6374A line.

Table 91a

Coronal observations at Sacramento Peak, New Mexico (5502A), east limb

Date GCT	Degrees north of the solar equator																			Degrees south of the solar equator																			
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
1954																																							
May 2.6	-	-	-	-	-	-	-	2	3	4	5	4	3	3	3	3	4	4	8	9	11	10	8	4	3	2	2	2	3	3	2	-	-	-	-	-	-	-	
3.7a	-	-	-	-	-	-	-	-	-	3	3	2	3	3	3	3	3	2	3	5	5	4	3	3	2	3	3	3	3	2	-	-	-	-	-	-	-	-	
4.7a	-	-	-	-	-	-	-	3	3	3	4	3	3	3	3	3	4	4	4	4	4	5	4	4	3	3	4	4	3	2	-	-	-	-	-	-	-	-	
5.7a	-	-	-	-	-	-	-	2	2	3	4	3	2	3	2	2	3	2	2	3	4	3	4	5	4	4	2	2	3	3	2	-	-	-	-	-	-	-	
6.7a	-	-	-	-	-	-	-	2	2	3	3	3	2	2	3	3	2	2	2	3	3	4	3	3	2	2	3	3	2	2	-	-	-	-	-	-	-	-	
8.7a	-	-	-	-	-	-	-	2	3	3	4	4	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
10.7a	-	-	-	-	-	-	-	2	2	3	3	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
11.8a	-	-	-	-	-	-	-	-	-	2	2	3	2	-	-	-	-	-	-	-	-	-	-	-	-	3	3	-	-	-	-	-	-	-	-	-	-	-	
12.8a	-	-	-	-	-	-	-	2	2	2	3	3	3	2	3	2	2	2	2	3	3	2	2	2	2	3	2	-	-	-	-	-	-	-	-	-	-	-	
19.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
23.6	-	-	-	-	-	-	-	-	-	2	2	3	2	2	2	2	2	2	2	3	2	2	-	-	2	2	2	3	3	2	-	-	-	-	-	-	-	-	
25.7	-	-	-	-	-	-	-	2	3	2	2	3	2	2	3	3	2	3	3	2	2	3	2	2	2	-	2	3	3	2	3	-	-	-	-	-	-	-	
26.7a	-	-	-	-	-	-	-	3	3	2	2	3	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
28.7	-	-	-	-	-	-	2	3	3	3	3	4	4	3	2	-	-	-	2	4	3	4	5	4	3	2	2	2	-	-	-	-	-	-	-	-	-	-	
30.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	2	3	3	3	4	3	2	2	2	-	-	-	-	-	-	-	-	-	-	
31.7	-	-	-	-	-	-	-	2	2	2	2	3	3	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

Table 92a

Coronal observations at Sacramento Peak, New Mexico (6574A), east limb

Date GCT	Degrees north of the solar equator																			Degrees south of the solar equator																			
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
1954																																							
May 2.6	3	4	3	3	4	2	-	-	2	2	3	4	5	6	5	8	10	13	11	9	8	7	5	4	5	8	10	9	5	3	-	2	3	3	3	3	4		
3.7a	2	3	3	2	2	2	2	3	3	3	3	3	4	5	5	5	6	6	7	5	4	4	4	5	4	4	6	5	3	3	-	2	2	3	3	4			
4.7a	3	2	3	2	3	-	2	2	3	3	2	2	3	3	5	5	6	7	8	9	7	7	5	5	6	5	4	3	2	-	3	2	-	2	2	3			
5.7a	2	3	3	2	2	2	2	3	3	3	2	3	2	4	5	4	6	8	7	6	5	5	4	4	3	4	4	3	3	2	2	2	2	2	3	3			
6.7a	2	3	2	2	-	-	-	2	3	-	2	3	4	4	5	6	7	8	7	7	8	6	5	6	5	4	4	5	2	2	-	2	-	-	3	3			
8.7a	2	2	2	-	-	-	-	-	-	-	-	-	3	3	3	4	4	4	4	5	4	4	4	3	2	3	5	3	3	2	2	2	-	-	-	-			
10.7a	2	-	-	-	-	-	-	-	-	-	-	2	3	3	4	5	5	6	7	6	5	5	4	5	5	5	3	4	2	2	2	2	2	2	2	2			
11.8a	-	2	2	-	2	3	2	-	2	3	6	5	6	4	5	7	7	8	9	8	7	6	5	6	5	5	6	-	-	-	-	-	-	-	-	-			
12.8a	2	3	3	3	3	-	-	2	3	2	3	3	4	5	4	5	7	8	9	8	6	5	5	6	5	4	3	2	2	2	3	3	2	2	2	3	3		
19.7	2	2	3	2	-	2	2	-	2	3	4	4	5	4	5	5	8	7	8	6	5	4	4	4	5	4	3	2	2	-	-	-	2	3	3	3			
23.6	2	2	-	2	-	2	2	2	2	2	3	2	3	3	4	5	4	4	4	4	4	3	4	5	4	7	5	4	4	3	2	2	3	2	3	3			
25.7	2	3	2	2	2	3	2	2	2	3	3	2	-	3	5	6	5	6	7	8	10	9	8	5	5	5	6	4	3	2	-	2	2	3	2	3			
26.7a	-	-	-	-	-	-	-	-	-	2	2	3	3	3	3	4	5	6	8	7	6	5	4	4	3	3	4	5	3	-	-	-	2	3	4	4	3		
28.7	4	2	2	2	3	2	2	-	2	3	3	5	7	8	7	8	8	6	5	5	6	4	4	4	3	4	5	2	3	2	-	2	3	3	3	3			
30.7a	-	-	-	-	-	-	-	-	-	2	2	2	3	4	5	4	5	4	5	4	3	3	4	5	4	3	3	2	2	-	-	-	-	-	-	-			
31.7	2	3	2	2	3	2	2	-	-	3	4	6	5	6	8	9	11	12	11	10	8	9	9	8	7	8	5	2	2	-	-	2	-	2	3	4			

Table 93a

Coronal observations at Sacramento Peak, New Mexico (6702A), east limb

The 1954 coronal line observations were made on an 8" telescope with a 10" filter.

Table 91b

Coronal observations at Sacramento Peak, New Mexico (530JA), west limb

Date GMT	Degrees south of the solar equator															Degrees north of the solar equator																								
	0	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	0	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90		
1954																																								
May 2.6	-	-	-	-	-	-	-	2	3	3	4	3	3	3	2	2	-	-	-	-	-	-	-	-	-	2	2	2	3	4	3	2	-	-	-	-	-	-	-	-
3.7	-	-	-	-	-	-	-	-	2	3	3	2	2	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	3	3	3	3	-	-	-	-	-	
4.7a	-	-	-	-	-	-	-	2	2	2	3	3	3	5	4	4	3	2	3	3	3	4	3	3	3	2	2	2	3	2	-	-	-	-	-	-	-	-	-	
5.7a	-	-	-	-	-	-	-	2	2	2	3	3	2	3	3	3	3	2	3	3	3	2	3	3	2	2	2	2	3	-	-	-	-	-	-	-	-	-	-	
6.7a	-	-	-	-	-	-	-	-	2	3	3	2	3	2	3	3	2	2	2	3	3	2	2	3	3	2	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	2	3	4	4	3	-	-	-	-	-	-	-	-	
10.7a	-	-	-	-	-	-	-	-	2	2	3	2	2	-	-	-	-	-	-	-	-	-	-	2	2	2	3	2	-	-	-	-	-	-	-	-	-	-	-	
11.8a	-	-	-	-	-	-	-	-	-	-	-	2	2	2	2	3	3	2	2	2	3	3	4	3	3	2	3	3	2	2	-	-	-	-	-	-	-	-	-	
12.8	-	-	-	-	-	-	-	-	-	2	3	3	2	3	2	2	2	-	-	-	-	-	2	2	-	-	2	3	2	-	-	-	-	-	-	-	-	-	-	
19.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	3	2	3	2	-	-	-	-	-	-	-	-	
23.6	-	-	-	-	-	-	-	-	-	2	2	2	-	-	2	3	4	3	3	2	2	2	2	2	-	-	3	3	2	-	-	-	-	-	-	-	-	-	-	
25.7	-	-	-	-	-	-	-	-	2	2	-	2	2	3	3	3	2	3	2	3	2	2	-	-	-	-	-	2	3	2	2	-	-	-	-	-	-	-	-	-
26.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
28.7	-	-	-	-	-	-	-	-	-	2	2	-	-	-	-	-	-	-	-	2	2	-	-	2	2	-	2	3	3	2	2	-	-	-	-	-	-	-	-	
30.7a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	2	3	3	3	2	2	2	-	-	-	-	-	-	-	-	-	-	-		
31.7	-	-	-	-	-	-	-	2	3	3	2	3	4	3	2	2	-	-	-	-	-	-	-	2	3	2	2	2	2	-	-	-	-	-	-	-	-	-	-	

Table 92b

Coronal observations at Sacramento Peak, New Mexico (6374A), west limb

Date GMT	Degrees south of the solar equator																			Degrees north of the solar equator																						
	90	85	80	75	70	65	60	55	50	45	40	35	30	25	20	15	10	5	0	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90					
1954																																										
May 2.6	4	3	3	2	3	2	-	2	-	3	2	3	6	7	8	7	7	7	10	7	8	10	9	8	12	11	8	3	2	2	-	-	2	3	4	3	3					
3.7	4	4	3	3	3	2	2	-	2	3	3	3	4	6	6	5	8	10	11	9	7	6	5	6	8	9	8	6	2	3	-	-	2	2	2	2	2					
4.7a	3	3	2	2	2	3	2	2	3	4	3	3	4	8	7	5	6	8	7	5	6	4	4	3	4	4	3	3	2	2	-	2	2	2	3	2	3					
5.7a	3	3	2	2	2	2	2	2	3	3	3	4	5	8	7	7	6	7	11	10	8	7	7	6	5	6	5	4	3	-	-	2	2	2	2	3	2					
6.7a	3	3	3	2	2	2	-	2	2	2	3	4	4	5	4	5	4	4	5	4	5	6	5	5	4	5	5	3	3	2	2	-	-	-	-	-	2	2				
8.7a	-	-	-	-	-	-	-	-	-	-	2	4	4	5	6	5	4	5	8	7	6	6	4	3	4	4	4	4	3	2	-	-	-	-	-	-	-	-	2			
10.7a	2	2	-	-	-	-	-	-	-	2	-	3	3	5	4	5	6	7	6	5	4	5	5	6	4	3	2	2	3	2	2	3	2	2	2	3	2	2				
11.8a	-	-	-	-	-	-	-	-	-	-	2	3	4	4	3	4	4	3	4	4	5	5	5	4	4	4	3	2	-	-	-	-	-	-	-	-	-	-				
12.8	3	2	3	2	2	2	3	2	2	3	2	2	3	4	5	5	4	6	8	7	7	7	7	6	5	5	4	-	2	2	-	-	-	2	3	2	3	2				
19.7	3	2	2	2	2	2	2	2	2	3	2	3	3	3	5	4	5	6	7	6	5	4	4	3	2	3	3	2	-	-	-	2	2	2	2	2	2					
23.6	3	2	2	2	2	2	2	2	2	2	2	2	3	3	4	4	3	4	5	6	5	6	5	6	8	7	8	7	7	3	2	-	2	2	-	3	3	3	2			
25.7	3	2	2	2	3	2	2	-	2	3	2	3	4	3	5	5	8	7	8	5	4	5	5	6	4	2	3	2	-	2	3	3	3	2	2	2	2	2				
26.7a	3	2	2	2	2	2	2	2	2	2	3	3	3	3	3	4	4	6	8	4	3	2	2	2	2	3	3	-	-	-	-	-	-	-	-	-	-	-				
28.7	3	2	2	2	-	-	-	-	-	-	2	3	3	4	5	6	7	11	10	9	8	5	4	4	4	5	4	4	3	4	-	-	-	-	-	2	3	2	4			
30.7a	-	-	-	-	-	-	-	-	-	2	2	3	3	5	4	5	8	7	6	5	4	4	7	8	7	4	5	2	2	-	-	-	-	-	-	-	-	-	-			
31.7	4	3	2	3	3	2	-	2	2	-	2	3	8	7	8	8	14	13	11	9	8	5	10	8	7	6	5	4	3	2	2	3	2	3	2	3	2	2				

Table 93b

Coronal observations at Sacramento Peak, New Mexico (6702A), west limb

The 6702A coronal line was not visible on any of the observation dates in May.

Table 94Zürich Provisional Relative Sunspot NumbersMay 1954

Date	R _Z *	Date	R _Z *
1	0	17	0
2	0	18	0
3	0	19	0
4	0	20	0
5	8	21	0
6	0	22	0
7	0	23	0
8	0	24	0
9	0	25	0
10	0	26	0
11	0	27	0
12	0	28	0
13	0	29	0
14	9	30	0
15	6	31	0
16	0	Mean:	0.7

*Dependent on observations at Zürich Observatory and its stations at Locarno and Arosa.

Table 95American Relative Sunspot NumbersApril 1954

Date	RA'	Date	RA'
1	0	17	1
2	0	18	2
3	0	19	0
4	0	20	0
5	0	21	0
6	0	22	0
7	2	23	0
8	5	24	0
9	2	25	0
10	0	26	0
11	0	27	0
12	0	28	0
13	0	29	0
14	1	30	0
15	3	Mean: 4.0	
16	3		

Table 96Solar Flares, April and May 1954

No solar flares were reported for the months of April and May.

Table 98

Sudden Ionosphere Disturbances Observed at Washington, D. C.

April and May 1954

No sudden ionosphere disturbances were observed during the months
of April and May.

Note: Observers are invited to send to the CRPL information
on times of beginning and end of sudden ionosphere disturbances
for publication as above. Address letters to the Central Radio
Propagation Laboratory, National Bureau of Standards, Boulder,
Colorado.

GRAPHS OF IONOSPHERIC DATA




Fig. 1. WASHINGTON, D. C.
38.7°N, 77.1°W

MAY 1954




Fig. 2. WASHINGTON, D. C.

MAY 1954




Fig. 3. FAIRBANKS, ALASKA
64.9°N, 147.8°W

MARCH 1954




Fig. 4. FAIRBANKS, ALASKA

MARCH 1954




Fig. 5. ANCHORAGE, ALASKA
61.2°N, 149.9°W

MARCH 1954




Fig. 6. ANCHORAGE, ALASKA

MARCH 1954




Fig. 7. ADAK, ALASKA
51.9°N, 176.6°W

MARCH 1954




Fig. 8. ADAK, ALASKA

MARCH 1954




Fig. 9. GRAZ, AUSTRIA
47.1°N, 15.5°E

MARCH 1954




Fig. 10. GRAZ, AUSTRIA

MARCH 1954




Fig. 11. SAN FRANCISCO, CALIFORNIA
37.4°N, 122.2°W

MARCH 1954




Fig. 12. SAN FRANCISCO, CALIFORNIA MARCH 1954




Fig. 13. WHITE SANDS, NEW MEXICO
32.3°N, 106.5°W MARCH 1954




Fig. 14. WHITE SANDS, NEW MEXICO MARCH 1954




Fig. 15. FORMOSA, CHINA
25.0°N, 121.5°E MARCH 1954




Fig. 16. FORMOSA, CHINA MARCH 1954




Fig. 17. MAUI, HAWAII
20.8°N, 156.5°W

MARCH 1954




Fig. 18. MAUI, HAWAII

MARCH 1954




Fig. 19. PUERTO RICO, W.I.
18.5°N, 67.2°W

MARCH 1954








Fig. 20. PUERTO RICO, W.I.

MARCH 1954






Fig. 25. FAIRBANKS, ALASKA

64.9°N, 147.8°W

FEBRUARY 1954




Fig. 26. FAIRBANKS, ALASKA

FEBRUARY 1954




Fig. 27. ADAK, ALASKA

51.9°N, 176.6°W

FEBRUARY 1954




Fig. 28. ADAK, ALASKA

FEBRUARY 1954




Fig. 29. GUAM I.

13.6°N, 144.9°E

FEBRUARY 1954




Fig. 30. GUAM I.

FEBRUARY 1954




Fig. 31. LEOPOLDVILLE, BELGIAN CONGO

4.3°S, 15.3°E

FEBRUARY 1954




Fig. 32. LEOPOLDVILLE, BELGIAN CONGO

FEBRUARY 1954




Fig. 33. JOHANNESBURG, UNION OF S. AFRICA
26.2°S, 28.1°E
FEBRUARY 1954




Fig. 34. JOHANNESBURG, UNION OF S. AFRICA




Fig. 35. CAPETOWN, UNION OF S. AFRICA
34.2°S, 18.3°E
FEBRUARY 1954




Fig. 36. CAPETOWN, UNION OF S. AFRICA




Fig. 37. RESOLUTE BAY, CANADA
74.7°N, 94.9°W JANUARY 1954




Fig. 38. RESOLUTE BAY, CANADA JANUARY 1954




Fig. 39. FAIRBANKS, ALASKA
64.9°N, 147.8°W JANUARY 1954




Fig. 40. FAIRBANKS, ALASKA JANUARY 1954




Fig. 41. BAKER LAKE, CANADA
64.3°N, 96.0°W JANUARY 1954




Fig. 42. BAKER LAKE, CANADA JANUARY 1954




Fig. 43. CHURCHILL, CANADA
58.8°N, 94.2°W JANUARY 1954




Fig. 44. CHURCHILL, CANADA JANUARY 1954




Fig. 45. FORT CHIMO, CANADA

58.1°N, 68.3°W

JANUARY 1954




Fig. 46. FORT CHIMO, CANADA

JANUARY 1954




Fig. 47. PRINCE RUPERT, CANADA

54.3°N, 130.3°W

JANUARY 1954




Fig. 48. PRINCE RUPERT, CANADA

JANUARY 1954




Fig. 49. WINNIPEG, CANADA
49.9°N, 97.4°W

JANUARY 1954




Fig. 50. WINNIPEG, CANADA

JANUARY 1954




Fig. 51. OTTAWA, CANADA
45.4°N, 75.9°W

JANUARY 1954




Fig. 52. OTTAWA, CANADA

JANUARY 1954




Fig. 53. JOHANNESBURG, UNION OF S. AFRICA
26.2°S, 28.1°E
JANUARY 1954




Fig. 54. JOHANNESBURG, UNION OF S. AFRICA




Fig. 55. CAPETOWN, UNION OF S. AFRICA
34.2°S, 18.3°E
JANUARY 1954




Fig. 56. CAPETOWN, UNION OF S. AFRICA




Fig. 57. BAKER LAKE, CANADA
64.3°N, 96.0°W

DECEMBER 1953




Fig. 58. BAKER LAKE, CANADA

DECEMBER 1953




Fig. 59. REYKJAVIK, ICELAND
64.1°N, 21.8°W

DECEMBER 1953




Fig. 60. REYKJAVIK, ICELAND

DECEMBER 1953




Fig. 61. WAKKANAI, JAPAN
45.4°N, 141.7°E

DECEMBER 1953




Fig. 62. WAKKANAI, JAPAN

DECEMBER 1953




Fig. 63. AKITA, JAPAN
39.7°N, 140.1°E

DECEMBER 1953




Fig. 64. AKITA, JAPAN

DECEMBER 1953




Fig. 65. TOKYO, JAPAN
35.7°N, 139.5°E

DECEMBER 1953




Fig. 66. TOKYO, JAPAN

DECEMBER 1953




Fig. 67. YAMAGAWA, JAPAN
31.2°N, 130.6°E

DECEMBER 1953




Fig. 68. YAMAGAWA, JAPAN

DECEMBER 1953




Fig. 69. CALCUTTA, INDIA

22.6°N, 88.4°E

DECEMBER 1953




Fig. 70. CALCUTTA, INDIA

DECEMBER 1953




Fig. 71. RAROTONGA I.

21.3°S, 159.8°W

DECEMBER 1953




Fig. 72. RAROTONGA I.

DECEMBER 1953




Fig. 73. INVERNESS, SCOTLAND
57.4°N, 4.2°W NOVEMBER 1953




Fig. 74. INVERNESS, SCOTLAND NOVEMBER 1953




Fig. 75. SLOUGH, ENGLAND
51.5°N, 0.6°W NOVEMBER 1953




Fig. 76. SLOUGH, ENGLAND NOVEMBER 1953




Fig. 77. DELHI, INDIA
28.6°N, 77.1°E

NOVEMBER 1953




Fig. 78. DELHI, INDIA

NOVEMBER 1953




Fig. 79. CALCUTTA, INDIA
22.6°N, 88.4°E

NOVEMBER 1953




Fig. 80. CALCUTTA, INDIA

NOVEMBER 1953




Fig. 81. BOMBAY, INDIA
19.0°N, 73.0°E NOVEMBER 1953




Fig. 82. BOMBAY, INDIA NOVEMBER 1953




Fig. 83. MADRAS, INDIA
13.0°N, 80.2°E NOVEMBER 1953




Fig. 84. MADRAS, INDIA NOVEMBER 1953




Fig. 85. TIRUCHY, INDIA
10.8°N, 78.8°E

NOVEMBER 1953




Fig. 86. TIRUCHY, INDIA

NOVEMBER 1953




Fig. 87. SINGAPORE, BRITISH MALAYA
1.3°N, 103.8°E

NOVEMBER 1953




Fig. 88. SINGAPORE, BRITISH MALAYA

NOVEMBER 1953




Fig. 89. RAROTONGA I.
21.3°S, 159.8°W

NOVEMBER 1953




Fig. 90. RAROTONGA I.

NOVEMBER 1953




Fig. 91. INVERNESS, SCOTLAND
57.4°N, 4.2°W

OCTOBER 1953




Fig. 92. INVERNESS, SCOTLAND

OCTOBER 1953




Fig. 93. SLOUGH, ENGLAND
51.5°N, 0.6°W

OCTOBER 1953




Fig. 94. SLOUGH, ENGLAND

OCTOBER 1953




Fig. 95. DELHI, INDIA
28.6°N, 77.1°E

OCTOBER 1953




Fig. 96. DELHI, INDIA

OCTOBER 1953




Fig. 97. CALCUTTA, INDIA
22.6°N, 88.4°E

OCTOBER 1953




Fig. 98. CALCUTTA, INDIA

OCTOBER 1953




Fig. 99. BOMBAY, INDIA
19.0°N, 73.0°E

OCTOBER 1953




Fig. 100. BOMBAY, INDIA

OCTOBER 1953




Fig. 101. KHARTOUM, SUDAN

15.6°N, 32.6°E

OCTOBER 1953




Fig. 102. KHARTOUM, SUDAN

OCTOBER 1953




Fig. 103. MADRAS, INDIA

13.0°N, 80.2°E

OCTOBER 1953




Fig. 104. MADRAS, INDIA

OCTOBER 1953




Fig. 105. TIRUCHY, INDIA
10.8°N, 78.8°E

OCTOBER 1953




Fig. 106. TIRUCHY, INDIA

OCTOBER 1953




Fig. 107. SINGAPORE, BRITISH MALAYA
1.3°N, 103.8°E

OCTOBER 1953




Fig. 108. SINGAPORE, BRITISH MALAYA

OCTOBER 1953




Fig. 109. NAIROBI, KENYA
1.3°S, 36.8°E

OCTOBER 1953




Fig. 110. NAIROBI, KENYA

OCTOBER 1953




Fig. 111. SAO PAULO, BRAZIL
23.5°S, 46.5°W

OCTOBER 1953







Fig. 112. SAO PAULO, BRAZIL


OCTOBER 1953



OCTOBER 1953



OCTOBER 1953






Fig. 117. DELHI, INDIA
28.6°N, 77.1°E

SEPTEMBER 1953




Fig. 118. DELHI, INDIA

SEPTEMBER 1953




Fig. 119. CALCUTTA, INDIA
22.6°N, 88.4°E

SEPTEMBER 1953




Fig. 120. CALCUTTA, INDIA

SEPTEMBER 1953




Fig.121. BOMBAY, INDIA
19.0° N, 73.0° E

SEPTEMBER 1953




Fig.122. BOMBAY, INDIA

SEPTEMBER 1953




Fig.123. MADRAS, INDIA
13.0° N, 80.2° E

SEPTEMBER 1953




Fig.124. MADRAS, INDIA

SEPTEMBER 1953




Fig. 125. TIRUCHY, INDIA
10.8°N, 78.8°E

SEPTEMBER 1953




Fig. 126. TIRUCHY, INDIA

SEPTEMBER 1953




Fig. 127. SAO PAULO, BRAZIL
23.5°S, 46.5°W

SEPTEMBER 1953




Fig. 128. SAO PAULO, BRAZIL

SEPTEMBER 1953




Fig. 129. FALKLAND IS.
51.7°S, 57.8°W SEPTEMBER 1953




Fig. 130. FALKLAND IS. SEPTEMBER 1953




Fig. 131. PORT LOCKROY
64.8°S, 63.5°W SEPTEMBER 1953




Fig. 132. PORT LOCKROY SEPTEMBER 1953




Fig. 133. DELHI, INDIA
28.6°N, 77.1°E

AUGUST 1953




Fig. 134. DELHI, INDIA

AUGUST 1953




Fig. 135. BOMBAY, INDIA
19.0°N, 73.0°E

AUGUST 1953




Fig. 136. BOMBAY, INDIA

AUGUST 1953




Fig. 137. MADRAS, INDIA
13.0°N, 80.2°E

AUGUST 1953




Fig. 138. MADRAS, INDIA

AUGUST 1953




Fig. 139. TIRUCHY, INDIA
10.8°N, 78.8°E

AUGUST 1953




Fig. 140. TIRUCHY, INDIA

AUGUST 1953




Fig. 141. SAO PAULO, BRAZIL
23.5°S, 46.5°W

AUGUST 1953




Fig. 142. SAO PAULO, BRAZIL

AUGUST 1953




Fig. 143. PORT LOCKROY
64.8°S, 63.5°W

AUGUST 1953




Fig. 144. PORT LOCKROY

AUGUST 1953

Index of Tables and Graphs of Ionospheric Data

in CRPL-F118

	<u>Table page</u>	<u>Figure page</u>
Adak, Alaska		
March 1954	12	50
February 1954	14	55
Akita, Japan		
December 1953	17	64
Anchorage, Alaska		
March 1954	12	50
Baker Lake, Canada		
January 1954	15	59
December 1953	16	63
Bombay, India		
November 1953	18	69
October 1953	20	73
September 1953	22	79
August 1953	23	82
Calcutta, India		
December 1953	17	66
November 1953	18	68
October 1953	20	73
September 1953	21	78
Capetown, Union of S. Africa		
February 1954	14	57
January 1954	16	62
Churchill, Canada		
January 1954	15	59
Delhi, India		
November 1953	18	68
October 1953	19	72
September 1953	21	78
August 1953	23	82
Fairbanks, Alaska		
March 1954	12	49
February 1954	14	55
January 1954	15	58
Falkland Is.		
October 1953	21	77
September 1953	22	81
Formosa, China		
March 1954	13	52
Fort Chimo, Canada		
January 1954	15	60
Graz, Austria		
March 1954	12	51

Index (CHPL-F118, continued)

	<u>Table page</u>	<u>Figure page</u>
Guam I.		
March 1954	13	54
February 1954	14	56
Inverness, Scotland		
November 1953	18	67
October 1953	19	71
Johannesburg, Union of S. Africa		
February 1954	14	57
January 1954	16	62
Khartoum, Sudan		
October 1953	20	74
Kiruna, Sweden		
February 1954	13	54
Leopoldville, Belgian Congo		
February 1954	14	56
Madras, India		
November 1953	18	69
October 1953	20	74
September 1953	22	79
August 1953	23	83
Maui, Hawaii		
March 1954	13	53
Nairobi, Kenya		
October 1953	21	76
Ottawa, Canada		
January 1954	16	61
Port Lockroy		
October 1953	21	77
September 1953	22	81
August 1953	23	84
Prince Rupert, Canada		
January 1954	15	60
Puerto Rico, W. I.		
March 1954	13	53
Rarotonga I.		
December 1953	17	66
November 1953	19	71
Resolute Bay, Canada		
January 1954	15	58
Reykjavik, Iceland		
December 1953	16	63
San Francisco, California		
March 1954	12	51

Index (CRPL-F118, concluded)

	<u>Table page</u>	<u>Figure page</u>
Sao Paulo, Brazil		
October 1953	21	76
September 1953	22	80
August 1953	23	84
Singapore, British Malaya		
November 1953	19	70
October 1953	20	75
Slough, England		
November 1953	18	67
October 1953	19	72
Tiruchy, India		
November 1953	19	70
October 1953	20	75
September 1953	22	80
August 1953	23	83
Tokyo, Japan		
December 1953	17	65
Wakkanai, Japan		
December 1953	17	64
Washington, D. C.		
May 1954	12	49
White Sands, New Mexico		
March 1954	13	52
Winnipeg, Canada		
January 1954	16	61
Yamagawa, Japan		
December 1953	17	65



CRPL Reports

[A detailed list of CRPL publications is available from the Central Radio Propagation Laboratory upon request]

Daily:

Radio disturbance forecasts, every half hour from broadcast stations WWV and WWVH of the National Bureau of Standards.

Telephoned and telegraphed reports of ionospheric, solar, geomagnetic, and radio propagation data.

Semiweekly:

CRPL—J. North Atlantic Radio Propagation Forecast (of days most likely to be disturbed during following month).

CRPL—Jp. North Pacific Radio Propagation Forecast (of days most likely to be disturbed during following month).

Semimonthly:

CRPL—Ja. Semimonthly Frequency Revision Factors For CRPL Basic Radio Propagation Prediction Reports.

Monthly:

CRPL—D. Basic Radio Propagation Predictions—Three months in advance. (Dept. of the Army, TB 11-499-, monthly supplements to TM 11-499; Dept. of the Navy, DNC 13 () series; Dept. of the Air Force, TO 16-1B-2 series.) On sale by Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C. Members of the Armed Forces should address cognizant military office.

CRPL—F. Ionospheric Data. Limited distribution. This publication is in general disseminated only to those individuals or scientific organizations which collaborate in the exchange of ionospheric, solar, geomagnetic or other radio propagation data or in exchange for copies of publications on radio, physics and geophysics for the CRPL library.

Circulars of the National Bureau of Standards pertaining to Radio Sky Wave Transmission:

NBS Circular 462. Ionospheric Radio Propagation.

NBS Circular 465. Instructions for the Use of Basic Radio Propagation Predictions.

These circulars are on sale by the Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C. Members of the Armed Forces should address the respective military office having cognizance of radio wave propagation.

The publications listed above may be obtained without charge from the Central Radio Propagation Laboratory, unless otherwise indicated.
